

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
11 July 2002 (11.07.2002)

PCT

(10) International Publication Number
WO 02/053596 A2(51) International Patent Classification⁷: C07K 16/28,
A61K 39/395, C12N 5/06, G01N 33/574, 33/577, 33/68,
A61P 35/00, C12N 15/13, A01K 67/027, A61K 48/00

(21) International Application Number: PCT/US01/51113

(22) International Filing Date:
20 December 2001 (20.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/259,927 5 January 2001 (05.01.2001) US(71) Applicants (for all designated States except US): **PFIZER INC.** [US/US]; 235 East 42nd Street, New York, NY 10017 (US). **ABGENIX, INC.** [US/US]; 7601 Dumbarton Circle, Fremont, CA 94555 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **COHEN, Bruce, D.** [US/US]; 41 Cardinal Road, East Lyme, CT 06333 (US). **BEEBE, Jean** [US/US]; 383 Forsyth Road, Salem, CT 06420 (US). **MILLER, Penelope, E.** [US/US]; 56 Hancock Drive, Mystic, CT 06355 (US). **MOYER, James, D.** [US/US]; 5 Jefferson Drive, East Lyme, CT 06333 (US). **CORVALAN, Jose, R.** [US/US]; 125 Williams lane, Foster City, CA 94404 (US). **GALLO, Michael** [US/CA]; 2650 Colwood Drive, North Vancouver, British Columbia V7R 2R1 (US).(74) Agents: **HALEY, James, F.** et al.; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,

[Continued on next page]

(54) Title: ANTIBODIES TO INSULIN-LIKE GROWTH FACTOR I RECEPTOR

(57) Abstract: The present invention relates to antibodies and antigen-binding portions thereof that specifically bind to insulin-like growth factor I receptor (IGF-IR), which is preferably human IGF-IR. The invention also relates to human anti-IGF-IR antibodies, including chimeric, bispecific, derivatized, single chain antibodies or portions of fusion proteins. The invention also relates to isolated heavy and light chain immunoglobulin molecules derived from anti-IGF-IR antibodies and nucleic acid molecules encoding such molecules. The present invention also relates to methods of making anti-IGF-IR antibodies, pharmaceutical compositions comprising these antibodies and methods of using the antibodies and compositions thereof for diagnosis and treatment. The invention also provides gene therapy methods using nucleic acid molecules encoding the heavy and/or light immunoglobulin molecules that comprise the human anti-IGF-IR antibodies. The invention also relates to gene therapy methods and transgenic animals comprising nucleic acid molecules of the present invention.

WO 02/053596 A2



- MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- Published:**
- without international search report and to be republished upon receipt of that report
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ANTIBODIES TO INSULIN-LIKE GROWTH FACTOR I RECEPTOR

This application claims the benefit of United States Provisional
5 Application 60/259,927, filed January 5, 2001.

BACKGROUND OF THE INVENTION

Insulin-like growth factor (IGF-I) is a 7.5-kD polypeptide that
circulates in plasma in high concentrations and is detectable in most tissues. IGF-I
stimulates cell differentiation and cell proliferation, and is required by most
10 mammalian cell types for sustained proliferation. These cell types include, among
others, human diploid fibroblasts, epithelial cells, smooth muscle cells, T
lymphocytes, neural cells, myeloid cells, chondrocytes, osteoblasts and bone marrow
stem cells. For a review of the wide variety of cell types for which IGF-I/IGF-I
receptor interaction mediates cell proliferation, see Goldring et al., Eukar. Gene
15 Express., 1:31-326 (1991).

The first step in the transduction pathway leading to IGF-I-stimulated
cellular proliferation or differentiation is binding of IGF-I or IGF-II (or insulin at
supraphysiological concentrations) to the IGF-I receptor. The IGF-I receptor is
composed of two types of subunits: an alpha subunit (a 130-135 kD protein that is
20 entirely extracellular and functions in ligand binding) and a beta subunit (a 95-kD
transmembrane protein, with transmembrane and cytoplasmic domains). The IGF-IR
belongs to the family of tyrosine kinase growth factor receptors (Ullrich et al., Cell
61: 203-212, 1990), and is structurally similar to the insulin receptor (Ullrich et al.,
EMBO J. 5: 2503-2512, 1986). The IGF-IR is initially synthesized as a single chain

proreceptor polypeptide which is processed by glycosylation, proteolytic cleavage, and covalent bonding to assemble into a mature 460-kD heterotetramer comprising two alpha-subunits and two beta-subunits. The beta subunit(s) possesses ligand-activated tyrosine kinase activity. This activity is implicated in the signaling pathways mediating ligand action which involve autophosphorylation of the beta-subunit and phosphorylation of IGF-IR substrates.

In vivo, serum levels of IGF-I are dependent upon the presence of pituitary growth hormone (GH). Although the liver is a major site of GH-dependent IGF-I synthesis, recent work indicates that the majority of normal tissues also produce IGF-I. A variety of neoplastic tissues may also produce IGF-I. Thus IGF-I may act as a regulator of normal and abnormal cellular proliferation via autocrine or paracrine, as well as endocrine mechanisms. IGF-I and IGF-II bind to IGF binding proteins (IGFBPs) *in vivo*. The availability of free IGF for interaction with the IGF-1R is modulated by the IGFBPs. For a review of IGFBPs and IGF-I, see Grimberg et al., J. Cell. Physiol. 183: 1-9, 2000.

There is considerable evidence for a role for IGF-I and/or IGF-IR in the maintenance of tumor cells *in vitro* and *in vivo*. IGF-IR levels are elevated in tumors of lung (Kaiser et al., J. Cancer Res. Clin Oncol. 119: 665-668, 1993; Moody et al., Life Sciences 52: 1161-1173, 1993; Macauley et al., Cancer Res., 50: 2511-2517, 1990), breast (Pollak et al., Cancer Lett. 38: 223-230, 1987; Foekens et al., Cancer Res. 49: 7002-7009, 1989; Cullen et al., Cancer Res. 49: 7002-7009, 1990; Arteaga et al., J. Clin. Invest. 84: 1418-1423, 1989), prostate and colon (Remaole-Bennet et al., J. Clin. Endocrinol. Metab. 75: 609-616, 1992; Guo et al., Gastroenterol. 102: 1101-1108, 1992). Deregulated expression of IGF-I in prostate epithelium leads to neoplasia in transgenic mice (DiGiovanni et al., Proc. Natl. Acad. Sci. USA 97: 3455-60, 2000). In addition, IGF-I appears to be an autocrine stimulator of human gliomas (Sandberg-Nordqvist et al., Cancer Res. 53: 2475-2478, 1993), while IGF-I stimulated the growth of fibrosarcomas that overexpressed IGF-IR (Butler et al., Cancer Res. 58: 3021-27, 1998). Further, individuals with "high normal" levels of IGF-I have an increased risk of common cancers compared to individuals with IGF-I levels in the "low normal" range (Rosen et al., Trends Endocrinol. Metab. 10: 136-41, 1999). Many of these tumor cell types respond to IGF-I with a proliferative signal in culture (Nakanishi et al., J. Clin. Invest. 82: 354-

359, 1988; Freed et al., J. Mol. Endocrinol. 3: 509-514, 1989), and autocrine or paracrine loops for proliferation *in vivo* have been postulated (LeRoith et al., Endocrine Revs. 16: 143-163, 1995; Yee et al., Mol. Endocrinol. 3: 509-514, 1989). For a review of the role IGF-I/IGF-I receptor interaction plays in the growth of a
5 variety of human tumors, see Macaulay, Br. J. Cancer, 65: 311-320, 1992.

Increased IGF-I levels are also correlated with several noncancerous pathological states, including acromegaly and gigantism (Barkan, Cleveland Clin. J. Med. 65: 343, 347-349, 1998), while abnormal IGF-I/IGF-I receptor function has been implicated in psoriasis (Wraight et al., Nat. Biotech. 18: 521-526, 2000),
10 atherosclerosis and smooth muscle restenosis of blood vessels following angioplasty (Bayes-Genis et al., Circ. Res. 86: 125-130, 2000). Increased IGF-I levels also can be a problem in diabetes or in complications thereof, such as microvascular proliferation (Smith et al., Nat. Med. 5: 1390-1395, 1999). Decreased IGF-I levels, which occur, *inter alia*, in cases when GH serum levels are decreased or when there is an
15 insensitivity or resistance to GH, is associated with such disorders as small stature (Laron, Paediatr. Drugs 1: 155-159, 1999), neuropathy, decrease in muscle mass and osteoporosis (Rosen et al., Trends Endocrinol. Metab. 10: 136-141, 1999).

Using antisense expression vectors or antisense oligonucleotides to the IGF-IR RNA, it has been shown that interference with IGF-IR leads to inhibition of
20 IGF-I-mediated or IGF-II-mediated cell growth (see, e.g., Wraight et al., Nat. Biotech. 18: 521-526, 2000). The antisense strategy was successful in inhibiting cellular proliferation in several normal cell types and in human tumor cell lines. Growth can also be inhibited using peptide analogues of IGF-I (Pietrzkowski et al., Cell Growth & Diff. 3: 199-205, 1992; and Pietrzkowski et al., Mol. Cell. Biol., 12: 3883-3889,
25 1992), or a vector expressing an antisense RNA to the IGF-I RNA (Trojan et al., Science 259: 94-97, 1992). In addition, antibodies to IGF-IR (Arteaga et al., Breast Canc. Res. Treatm., 22: 101-106, 1992; and Kalebic et al., Cancer Res. 54: 5531-5534, 1994), and dominant negative mutants of IGF-IR (Prager et al., Proc. Natl. Acad. Sci. U.S.A. 91: 2181-2185, 1994; Li et al., J. Biol. Chem., 269: 32558-32564,
30 1994 and Jiang et al., Oncogene 18: 6071-77, 1999), can reverse the transformed phenotype, inhibit tumorigenesis, and induce loss of the metastatic phenotype.

IGF-I is also important in the regulation of apoptosis. Apoptosis, which is programmed cell death, is involved in a wide variety of developmental

processes, including immune and nervous system maturation. In addition to its role in development, apoptosis also has been implicated as an important cellular safeguard against tumorigenesis (Williams, *Cell* 65: 1097-1098, 1991; Lane, *Nature* 362: 786-787, 1993). Suppression of the apoptotic program, by a variety of genetic lesions, may contribute to the development and progression of malignancies.

IGF-I protects from apoptosis induced by cytokine withdrawal in IL-3-dependent hemopoietic cells (Rodriguez-Tarduchy, G. et al., *J. Immunol.* 149: 535-540, 1992), and from serum withdrawal in Rat-1/mycER cells (Harrington, E., et al., *EMBO J.* 13: 3286-3295, 1994). The anti-apoptotic function of IGF-I is important in the post-commitment stage of the cell cycle and also in cells blocked in cell cycle progression by etoposide or thymidine. The demonstration that *c-myc* driven fibroblasts are dependent on IGF-I for their survival suggests that there is an important role for the IGF-IR in the maintenance of tumor cells by specifically inhibiting apoptosis, a role distinct from the proliferative effects of IGF-I or IGF-IR. This would be similar to a role thought to be played by other anti-apoptotic genes such as *bcl-2* in promoting tumor survival (McDonnell et al., *Cell* 57: 79-88, 1989; Hockenberry et al., *Nature* 348: 334-336, 1990).

The protective effects of IGF-I on apoptosis are dependent upon having IGF-IR present on cells to interact with IGF-I (Resnicoff et al., *Cancer Res.* 55: 3739-3741, 1995). Support for an anti-apoptotic function of IGF-IR in the maintenance of tumor cells was also provided by a study using antisense oligonucleotides to the IGF-IR that identified a quantitative relationship between IGF-IR levels, the extent of apoptosis and the tumorigenic potential of a rat syngeneic tumor (Rescinoff et al., *Cancer Res.* 55: 3739-3741, 1995). An overexpressed IGF-1R has been found to protect tumor cells *in vitro* from etoposide-induced apoptosis (Sell et al., *Cancer Res.* 55: 303-306, 1995) and, even more dramatically, that a decrease in IGF-1R levels below wild type levels caused massive apoptosis of tumor cells *in vivo* (Resnicoff et al., *Cancer Res.* 55: 2463-2469, 1995).

Potential strategies for inducing apoptosis or for inhibiting cell proliferation associated with increased IGF-I, increased IGF-II and/or increased IGF-IR receptor levels include suppressing IGF-I levels or IGF-II levels or preventing the binding of IGF-I to the IGF-IR. For example, the long acting somatostatin analogue octreotide has been employed to reduce IGF synthesis and/or secretion. Soluble IGF-

IR has been used to induce apoptosis in tumor cells *in vivo* and inhibit tumorigenesis in an experimental animal system (D'Ambrosio et al., Cancer Res. 56: 4013-20, 1996). In addition, IGF-IR antisense oligonucleotides, peptide analogues of IGF-I, and antibodies to IGF-IR have been used to decrease IGF-I or IGF-IR expression (see
5 *supra*). However, none of these compounds has been suitable for long-term administration to human patients. In addition, although IGF-I has been administered to patients for treatment of short stature, osteoporosis, decreased muscle mass, neuropathy or diabetes, the binding of IGF-I to IGFBPs has often made treatment with IGF-I difficult or ineffective.

10 Accordingly, in view of the roles that IGF-I and IGF-IR have in such disorders as cancer and other proliferative disorders when IGF-I and/or IGF-IR are overexpressed, and the roles that too little IGF-I and IGF-IR have in disorders such as short stature and frailty when either IGF-I and/or IGF-IR are underexpressed, it would be desirable to generate antibodies to IGF-IR that could be used to either inhibit or
15 stimulate IGF-IR. Although anti-IGF-IR antibodies have been reported to have been found in certain patients with autoimmune diseases, none of these antibodies has been purified and none has been shown to be suitable for inhibiting IGF-I activity for diagnostic or clinical procedures. See, e.g., Thompson et al., Pediat. Res. 32: 455-459, 1988; Tappy et al., Diabetes 37: 1708-1714, 1988; Weightman et al.,
20 Autoimmunity 16:251-257, 1993; Drexhage et al., Nether. J. of Med. 45:285-293, 1994. Thus, it would be desirable to obtain high-affinity human anti-IGF-IR antibodies that could be used to treat diseases in humans.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figs. 1A-1C show alignments of the nucleotide sequences of the light chain variable regions from six human anti-IGF-IR antibodies to each other and to germline sequences. Fig. 1A shows the alignment of the nucleotide sequences of the variable region of the light chain (VL) of antibodies 2.12.1 (SEQ ID NO: 1) 2.13.2 (SEQ ID NO: 5), 2.14.3 (SEQ ID NO: 9) and 4.9.2 (SEQ ID NO: 13) to each other and to the germline V κ A30 sequence (SEQ ID NO: 39). Fig. 1B shows the
30 alignment of the nucleotide sequence of VL of antibody 4.17.3 (SEQ ID NO: 17) to the germline V κ O12 sequence (SEQ ID NO: 41). Fig. 1C shows the alignment of the nucleotide sequence of VL of antibody 6.1.1 (SEQ ID NO: 21) to the germline V κ

A27 sequence (SEQ ID NO: 37). The alignments also show the CDR regions of the VL from each antibody. The consensus sequences for Figs. 1A-1C are shown in SEQ ID NOS: 53-55, respectively.

5 Figs. 2A-2D show alignments of the nucleotide sequences of the heavy chain variable regions from six human anti-IGF-IR antibodies to each other and to germline sequences. Fig. 2A shows the alignment of the nucleotide sequence of the VH of antibody 2.12.1 (SEQ ID NO: 3) to the germline VH DP-35 sequence (SEQ ID NO: 29). Fig. 2B shows the alignment of the nucleotide sequence of the VH of antibody 2.14.3 (SEQ ID NO: 11) to the germline VIV-4/4.35 sequence (SEQ ID NO: 43). Figs. 2C-1 and 2C-2 show the alignments of the nucleotide sequences of the VH of antibodies 2.13.2 (SEQ ID NO: 7), 4.9.2 (SEQ ID NO: 15) and 6.1.1 (SEQ ID NO: 23) to each other and to the germline VH DP-47 sequence (SEQ ID NO: 31). Fig. 2D shows the alignment of the nucleotide sequence of the VH of antibody 4.17.3 (SEQ ID NO: 19) to the germline VH DP-71 sequence (SEQ ID NO: 35). The alignment
10 also shows the CDR regions of the antibodies. The consensus sequences for Figs. 2A-2D are shown in SEQ ID NOS: 56-59, respectively.

Fig. 3 shows that anti-IGF-IR antibodies 2.13.2, 4.9.2 and 2.12.1 inhibit IGF-I binding to 3T3-IGF-IR cells.

Fig. 4 shows that anti-IGF-IR antibody 4.9.2 inhibits IGF-I-induced
20 receptor tyrosine phosphorylation (upper panel) and induces IGF-IR downregulation at the cell surface (lower panel).

Fig. 5 shows that anti-IGF-IR antibodies 2.13.2 and 4.9.2 reduce IGF-IR phosphotyrosine signal in 3T3-IGF-IR tumors.

Fig. 6 shows that anti-IGF-IR antibodies 2.13.2 and 4.9.2 reduce IGF-
25 IR in 3T3-IGF-IR tumors.

Fig. 7 shows that anti-IGF-IR antibody 2.13.2 inhibits 3T3-IGF-IR tumor growth *in vivo* alone (left panel) or in combination with adriamycin (right panel).

Fig. 8 shows the relationship between anti-IGF-IR antibody 2.13.2
30 serum levels and IGF-IR downregulation in 3T3-IGF-IR tumors.

Fig. 9 shows that multiple doses of anti-IGF-IR antibody 2.13.2 inhibit 3T3-IGF-IR tumor growth *in vivo* alone or in combination with adriamycin.

Fig. 10 shows that anti-IGF-IR antibody 2.13.2 inhibits large tumor growth *in vivo* in combination with adriamycin.

Fig. 11 shows that anti-IGF-IR antibody 2.13.2 inhibits Colo 205 tumor growth *in vivo* alone or in combination with 5-deoxyuridine (5-FU).

5 Fig. 12 shows that multiple doses of anti-IGF-IR antibody 2.13.2 inhibit Colo 205 tumor growth *in vivo* alone or in combination with 5-FU.

Fig. 13 shows that multiple doses of anti-IGF-IR antibody 2.13.2 inhibit MCF-7 tumor growth *in vivo* alone or in combination with taxol.

10 Fig. 14 shows that anti-IGF-IR antibody 2.13.2 inhibits MCF-7 tumor growth *in vivo* alone (left panel) or in combination with adriamycin (right panel).

Fig. 15 shows that multiple doses of anti-IGF-IR antibody 2.13.2 inhibit MCF-7 tumor growth *in vivo* alone or in combination with tamoxifen.

15 Fig. 16 shows that multiple doses of anti-IGF-IR antibody 2.13.2 inhibit A431 tumor growth *in vivo* alone or in combination with the epidermal growth factor receptor (EGF-R) tyrosine kinase inhibitor CP-358,774.

Fig. 17 shows a pharmacokinetic evaluation of a single intravenous injection of anti-IGF-IR antibody 2.13.2 in Cynomolgus monkeys.

Fig. 18 shows that the combination of anti-IGF-IR antibody 2.13.2 and adriamycin increases the downregulation of IGF-IR on 3T3-IGF-IR tumors *in vivo*.

20 Fig. 19A shows the number of mutations in different regions of the heavy and light chains of 2.13.2 and 2.12.1 compared to the germline sequences.

Figs. 19A-D show alignments of the amino acid sequences from the heavy and light chains of antibodies 2.13.2 and 2.12.1 with the germline sequences from which they are derived. Fig. 19B shows an alignment of the amino acid sequence of the heavy chain of antibody 2.13.2 (SEQ ID NO: 45) with that of germline sequence DP-47(3-23)/D6-19/JH6 (SEQ ID NO: 46). Fig. 19C shows an alignment of the amino acid sequence of the light chain of antibody 2.13.2 (SEQ ID NO: 47) with that of germline sequence A30/Jk2 (SEQ ID NO: 48). Fig. 19D shows an alignment of the amino acid sequence of the heavy chain of antibody 2.12.1 (SEQ ID NO: 49) with that of germline sequence DP-35(3-11)/D3-3/JH6 (SEQ ID NO: 50). Fig. 19E shows an alignment of the amino acid sequence of the light chain of antibody 2.12.1 (SEQ ID NO: 51) with that of germline sequence A30/Jk1 (SEQ ID NO: 52). For Figures 19B-E, the signal sequences are in italic, the CDRs are

25
30

underlined, the constant domains are bold, the framework (FR) mutations are highlighted with a plus sign (“+”) above the amino acid residue and CDR mutations are highlighted with an asterisk above the amino acid residue.

SUMMARY OF THE INVENTION

5 The present invention provides an isolated antibody or antigen-binding portion thereof that binds IGF-IR, preferably one that binds to primate and human IGF-IR, and more preferably one that is a human antibody. The invention provides an anti-IGF-IR antibody that inhibits the binding of IGF-I or IGF-II to IGF-IR, and also provides an anti-IGF-IR antibody that activates IGF-IR.

10 The invention provides a pharmaceutical composition comprising the antibody and a pharmaceutically acceptable carrier. The pharmaceutical composition may further comprise another component, such as an anti-tumor agent or an imaging reagent.

Diagnostic and therapeutic methods are also provided by the invention.

15 Diagnostic methods include a method for diagnosing the presence or location of an IGF-IR-expressing tissue using an anti-IGF-IR antibody. A therapeutic method comprises administering the antibody to a subject in need thereof, preferably in conjunction with administration of another therapeutic agent.

20 The invention provides an isolated cell line, such as a hybridoma, that produces an anti-IGF-IR antibody.

25 The invention also provides nucleic acid molecules encoding the heavy and/or light chain or antigen-binding portions thereof of an anti-IGF-IR antibody. The invention provides vectors and host cells comprising the nucleic acid molecules, as well as methods of recombinantly producing the polypeptides encoded by the nucleic acid molecules.

30 Non-human transgenic animals that express the heavy and/or light chain or antigen-binding portions thereof of an anti-IGF-IR antibody are also provided. The invention also provides a method for treating a subject in need thereof with an effective amount of a nucleic acid molecule encoding the heavy and/or light chain or antigen-binding portions thereof of an anti-IGF-IR antibody.

DETAILED DESCRIPTION OF THE INVENTION

Definitions and General Techniques

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992), and Harlow and Lane *Antibodies: A Laboratory Manual* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990), which are incorporated herein by reference. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclatures used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

The following terms, unless otherwise indicated, shall be understood to have the following meanings:

The term "polypeptide" encompasses native or artificial proteins, protein fragments and polypeptide analogs of a protein sequence. A polypeptide may be monomeric or polymeric.

The term "isolated protein" or "isolated polypeptide" is a protein or polypeptide that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) is free of other proteins from the same species (3) is expressed by a cell from a different species, or (4) does not occur in nature. Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be "isolated" from its naturally associated components. A protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art.

A protein or polypeptide is "substantially pure," "substantially homogeneous" or "substantially purified" when at least about 60 to 75% of a sample exhibits a single species of polypeptide. The polypeptide or protein may be monomeric or multimeric. A substantially pure polypeptide or protein will typically comprise about 50%, 60, 70%, 80% or 90% W/W of a protein sample, more usually about 95%, and preferably will be over 99% pure. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band upon staining the gel with a stain well known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art for purification.

The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence. Fragments typically are at least 5, 6, 8 or 10 amino acids long, preferably at least 14 amino acids long, more preferably at least 20 amino acids long, usually at least 50 amino acids long, even more preferably at least 70, 80, 90, 100, 150 or 200 amino acids long.

The term "polypeptide analog" as used herein refers to a polypeptide that is comprised of a segment of at least 25 amino acids that has substantial identity to a portion of an amino acid sequence and that has at least one of the following properties: (1) specific binding to IGF-IR under suitable binding conditions, (2) ability to block IGF-I or IGF-II binding to IGF-IR, or (3) ability to reduce IGF-IR cell surface expression or tyrosine phosphorylation *in vitro* or *in vivo*. Typically,

polypeptide analogs comprise a conservative amino acid substitution (or insertion or deletion) with respect to the naturally-occurring sequence. Analogs typically are at least 20 amino acids long, preferably at least 50, 60, 70, 80, 90, 100, 150 or 200 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

Preferred amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and (4) confer or modify other physicochemical or functional properties of such analogs. Analogs can include various muteins of a sequence other than the naturally-occurring peptide sequence. For example, single or multiple amino acid substitutions (preferably conservative amino acid substitutions) may be made in the naturally-occurring sequence (preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid substitution should not substantially change the structural characteristics of the parent sequence (e.g., a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al. *Nature* 354:105 (1991), which are each incorporated herein by reference.

Non-peptide analogs are commonly used in the pharmaceutical industry as drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, *J. Adv. Drug Res.* 15:29 (1986); Veber and Freidinger *TINS* p.392 (1985); and Evans et al. *J. Med. Chem.* 30:1229 (1987), which are incorporated herein by reference. Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a desired biochemical property or pharmacological

activity), such as a human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: --CH₂NH--, --CH₂S--, --CH₂-CH₂--, --CH=CH--(cis and trans), --COCH₂--, --CH(OH)CH₂--, and --CH₂SO--, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may also be used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.* 61:387 (1992), incorporated herein by reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

An "immunoglobulin" is a tetrameric molecule. In a naturally-occurring immunoglobulin, each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Human light chains are classified as κ and λ light chains. Heavy chains are classified as μ , Δ , γ , α , or ϵ , and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. *See generally, Fundamental Immunology* Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes). The variable regions of each light/heavy chain pair form the antibody binding site such that an intact immunoglobulin has two binding sites.

Immunoglobulin chains exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N-terminus to C-terminus, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of

Kabat *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk *J. Mol. Biol.* 196:901-917 (1987); Chothia et al. *Nature* 342:878-883 (1989).

An "antibody" refers to an intact immunoglobulin or to an antigen-binding portion thereof that competes with the intact antibody for specific binding. Antigen-binding portions may be produced by recombinant DNA techniques or by enzymatic or chemical cleavage of intact antibodies. Antigen-binding portions include, *inter alia*, Fab, Fab', F(ab')₂, Fv, dAb, and complementarity determining region (CDR) fragments, single-chain antibodies (scFv), chimeric antibodies, diabodies and polypeptides that contain at least a portion of an immunoglobulin that is sufficient to confer specific antigen binding to the polypeptide.

As used herein, an antibody that is referred to as, e.g., 2.12.1, 2.13.2, 2.14.3, 4.9.2, 4.17.3 and 6.1.1, is an antibody that is derived from the hybridoma of the same name. For example, antibody 2.12.1 is derived from hybridoma 2.12.1.

An Fab fragment is a monovalent fragment consisting of the VL, VH, CL and CH I domains; a F(ab')₂ fragment is a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consists of the VH and CH1 domains; an Fv fragment consists of the VL and VH domains of a single arm of an antibody; and a dAb fragment (Ward et al., *Nature* 341:544-546, 1989) consists of a VH domain.

A single-chain antibody (scFv) is an antibody in which a VL and VH regions are paired to form a monovalent molecules via a synthetic linker that enables them to be made as a single protein chain (Bird et al., *Science* 242:423-426, 1988 and Huston et al., *Proc. Natl. Acad. Sci. USA* 85:5879-5883, 1988). Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see e.g., Holliger, P., et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448, 1993, and Poljak, R. J., et al., *Structure* 2:1121-1123, 1994). One or more CDRs may be incorporated into a molecule either covalently or noncovalently to make it an immunoadhesin. An immunoadhesin may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the

CDR(s) noncovalently. The CDRs permit the immunoadhesin to specifically bind to a particular antigen of interest.

An antibody may have one or more binding sites. If there is more than one binding site, the binding sites may be identical to one another or may be different. For instance, a naturally-occurring immunoglobulin has two identical binding sites, a single-chain antibody or Fab fragment has one binding site, while a "bispecific" or "bifunctional" antibody has two different binding sites.

An "isolated antibody" is an antibody that (1) is not associated with naturally-associated components, including other naturally-associated antibodies, that accompany it in its native state, (2) is free of other proteins from the same species, (3) is expressed by a cell from a different species, or (4) does not occur in nature. Examples of isolated antibodies include an anti-IGF-IR antibody that has been affinity purified using IGF-IR is an isolated antibody, an anti-IGF-IR antibody that has been synthesized by a hybridoma or other cell line *in vitro*, and a human anti-IGF-IR antibody derived from a transgenic mouse.

The term "human antibody" includes all antibodies that have one or more variable and constant regions derived from human immunoglobulin sequences. In a preferred embodiment, all of the variable and constant domains are derived from human immunoglobulin sequences (a fully human antibody). These antibodies may be prepared in a variety of ways, as described below.

A humanized antibody is an antibody that is derived from a non-human species, in which certain amino acids in the framework and constant domains of the heavy and light chains have been mutated so as to avoid or abrogate an immune response in humans. Alternatively, a humanized antibody may be produced by fusing the constant domains from a human antibody to the variable domains of a non-human species. Examples of how to make humanized antibodies may be found in United States Patent Nos. 6,054,297, 5,886,152 and 5,877,293.

The term "chimeric antibody" refers to an antibody that contains one or more regions from one antibody and one or more regions from one or more other antibodies. In a preferred embodiment, one or more of the CDRs are derived from a human anti-IGF-IR antibody. In a more preferred embodiment, all of the CDRs are derived from a human anti-IGF-IR antibody. In another preferred embodiment, the CDRs from more than one human anti-IGF-IR antibodies are mixed and matched in a

chimeric antibody. For instance, a chimeric antibody may comprise a CDR1 from the light chain of a first human anti-IGF-IR antibody may be combined with CDR2 and CDR3 from the light chain of a second human anti-IGF-IR antibody, and the CDRs from the heavy chain may be derived from a third anti-IGF-IR antibody. Further, the framework regions may be derived from one of the same anti-IGF-IR antibodies, from one or more different antibodies, such as a human antibody, or from a humanized antibody.

A "neutralizing antibody" or "an inhibitory antibody" is an antibody that inhibits the binding of IGF-IR to IGF-I when an excess of the anti-IGF-IR antibody reduces the amount of IGF-I bound to IGF-IR by at least about 20%. In a preferred embodiment, the antibody reduces the amount of IGF-I bound to IGF-IR by at least 40%, more preferably 60%, even more preferably 80%, or even more preferably 85%. The binding reduction may be measured by any means known to one of ordinary skill in the art, for example, as measured in an *in vitro* competitive binding assay. An example of measuring the reduction in binding of IGF-I to IGF-IR is presented below in Example IV.

An "activating antibody" is an antibody that activates IGF-IR by at least about 20% when added to a cell, tissue or organism expressing IGF-IR. In a preferred embodiment, the antibody activates IGF-IR activity by at least 40%, more preferably 60%, even more preferably 80%, or even more preferably 85%. In a more preferred embodiment, the activating antibody is added in the presence of IGF-I or IGF-II. In another preferred embodiment, the activity of the activating antibody is measured by determining the amount of tyrosine autophosphorylation of IGF-IR.

Fragments or analogs of antibodies can be readily prepared by those of ordinary skill in the art following the teachings of this specification. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Preferably, computerized comparison methods are used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. Bowie et al. *Science* 253:164 (1991).

The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore system (Pharmacia Biosensor AB, Uppsala, Sweden and
5 Piscataway, N.J.). For further descriptions, see Jonsson, U., et al. (1993) *Ann. Biol. Clin.* 51:19-26; Jonsson, U., et al. (1991) *Biotechniques* 11:620-627; Johnsson, B., et al. (1995) *J. Mol. Recognit.* 8:125-131; and Johnson, B., et al. (1991) *Anal. Biochem.* 198:268-277.

The term " K_{off} " refers to the off rate constant for dissociation of an
10 antibody from the antibody/antigen complex.

The term " K_d " refers to the dissociation constant of a particular antibody-antigen interaction.

The term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor. Epitopic determinants
15 usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. An antibody is said to specifically bind an antigen when the dissociation constant is $\leq 1 \mu\text{M}$, preferably $\leq 100 \text{ nM}$ and most preferably $\leq 10 \text{ nM}$.

20 As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See *Immunology - A Synthesis* (2nd Edition, E.S. Golub and D.R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional
25 amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O⁻-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, s-N-
30 methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the lefthand direction is the amino terminal direction and the righthand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

The term "polynucleotide" as referred to herein means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

5 The term "isolated polynucleotide" as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the "isolated polynucleotide" (1) is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or
10 (3) does not occur in nature as part of a larger sequence.

 The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. Preferably
15 oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, e.g. for probes; although oligonucleotides may be double stranded, e.g. for use in the construction of a gene mutant. Oligonucleotides of the invention can be either sense or antisense oligonucleotides.

20 The term "naturally occurring nucleotides" referred to herein includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" referred to herein includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate,
25 phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See e.g., LaPlanche et al. *Nucl. Acids Res.* 14:9081 (1986); Stec et al. *J. Am. Chem. Soc.* 106:6077 (1984); Stein et al. *Nucl. Acids Res.* 16:3209 (1988); Zon et al. *Anti-Cancer Drug Design* 6:539 (1991); Zon et al. *Oligonucleotides and Analogues: A Practical Approach*, pp. 87-108 (F. Eckstein, Ed., Oxford University
30 Press, Oxford England (1991)); Stec et al. U.S. Patent No. 5,151,510; Uhlmann and Peyman *Chemical Reviews* 90:543 (1990), the disclosures of which are hereby incorporated by reference. An oligonucleotide can include a label for detection, if desired.

"Operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in *trans* or at a distance to control the gene of interest. The term "expression control sequence" as used herein refers to polynucleotide sequences which are necessary to effect the expression and processing of coding sequences to which they are ligated. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

The term "vector", as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid"

and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

5 The term "recombinant host cell" (or simply "host cell"), as used herein, is intended to refer to a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or
10 environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein.

 The term "selectively hybridize" referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof in accordance with the invention selectively hybridize to nucleic acid strands under
15 hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. "High stringency" or "highly stringent" conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. An example of "high stringency" or "highly stringent" conditions is a method of incubating a polynucleotide with another polynucleotide,
20 wherein one polynucleotide may be affixed to a solid surface such as a membrane, in a hybridization buffer of 6X SSPE or SSC, 50% formamide, 5X Denhardt's reagent, 0.5% SDS, 100 µg/ml denatured, fragmented salmon sperm DNA at a hybridization temperature of 42°C for 12-16 hours, followed by twice washing at 55°C using a wash buffer of 1X SSC, 0.5% SDS. See also Sambrook et al., *supra*, pp. 9.50-9.55.

25 The term "percent sequence identity" in the context of nucleic acid sequences refers to the residues in two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over a stretch of at least about nine nucleotides, usually at least about 18 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides,
30 more typically at least about 32 nucleotides, and preferably at least about 36, 48 or more nucleotides. There are a number of different algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using FASTA, Gap or Bestfit, which are programs in

Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wisconsin. FASTA, which includes, *e.g.*, the programs FASTA2 and FASTA3, provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, *Methods Enzymol.* 183: 63-98
5 (1990); Pearson, *Methods Mol. Biol.* 132: 185-219 (2000); Pearson, *Methods Enzymol.* 266: 227-258 (1996); Pearson, *J. Mol. Biol.* 276: 71-84 (1998); herein incorporated by reference). Unless otherwise specified, default parameters for a particular program or algorithm are used. For instance, percent sequence identity between nucleic acid sequences can be determined using FASTA with its default
10 parameters (a word size of 6 and the NOPAM factor for the scoring matrix) or using Gap with its default parameters as provided in GCG Version 6.1, herein incorporated by reference.

A reference to a nucleic acid sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid molecule having a
15 particular sequence should be understood to encompass its complementary strand, with its complementary sequence.

In the molecular biology art, researchers use the terms "percent sequence identity", "percent sequence similarity" and "percent sequence homology" interchangeably. In this application, these terms shall have the same meaning with
20 respect to nucleic acid sequences only.

The term "substantial similarity" or "substantial sequence similarity," when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about
25 85%, preferably at least about 90%, and more preferably at least about 95%, 96%, 97%, 98% or 99% of the nucleotide bases, as measured by any well-known algorithm of sequence identity, such as FASTA, BLAST or Gap, as discussed above.

As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or
30 BESTFIT using default gap weights, share at least 75% or 80% sequence identity, preferably at least 90% or 95% sequence identity, even more preferably at least 98% or 99% sequence identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. A "conservative amino acid substitution" is

one in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity or degree of similarity may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well-known to those of skill in the art. See, e.g., Pearson, *Methods Mol. Biol.* 24: 307-31 (1994), herein incorporated by reference. Examples of groups of amino acids that have side chains with similar chemical properties include 1) aliphatic side chains: glycine, alanine, valine, leucine and isoleucine; 2) aliphatic-hydroxyl side chains: serine and threonine; 3) amide-containing side chains: asparagine and glutamine; 4) aromatic side chains: phenylalanine, tyrosine, and tryptophan; 5) basic side chains: lysine, arginine, and histidine; and 6) sulfur-containing side chains are cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, glutamate-aspartate, and asparagine-glutamine.

Alternatively, a conservative replacement is any change having a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet et al., *Science* 256: 1443-45 (1992), herein incorporated by reference. A "moderately conservative" replacement is any change having a nonnegative value in the PAM250 log-likelihood matrix.

Sequence similarity for polypeptides, which is also referred to as sequence identity, is typically measured using sequence analysis software. Protein analysis software matches similar sequences using measures of similarity assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, GCG contains programs such as "Gap" and "Bestfit" which can be used with default parameters to determine sequence homology or sequence identity between closely related polypeptides, such as homologous polypeptides from different species of organisms or between a wild type protein and a mutein thereof. See, e.g., GCG Version 6.1. Polypeptide sequences also can be compared using FASTA using default or recommended parameters, a program in GCG Version 6.1. FASTA (e.g., FASTA2 and FASTA3) provides alignments and

percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson (1990); Pearson (2000)). Another preferred algorithm when comparing a sequence of the invention to a database containing a large number of sequences from different organisms is the computer program BLAST, especially
5 blastp or tblastn, using default parameters. See, e.g., Altschul et al., J. Mol. Biol. 215: 403-410 (1990); Altschul et al., Nucleic Acids Res. 25:3389-402 (1997); herein incorporated by reference.

The length of polypeptide sequences compared for homology will generally be at least about 16 amino acid residues, usually at least about 20 residues,
10 more usually at least about 24 residues, typically at least about 28 residues, and preferably more than about 35 residues. When searching a database containing sequences from a large number of different organisms, it is preferable to compare amino acid sequences.

As used herein, the terms "label" or "labeled" refers to incorporation of
15 another molecule in the antibody. In one embodiment, the label is a detectable marker, e.g., incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). In another embodiment, the label or marker can
20 be therapeutic, e.g., a drug conjugate or toxin. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish
25 peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent markers, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags), magnetic agents, such as gadolinium chelates, toxins such as pertussis toxin, taxol, cytochalasin B, gramicidin D, ethidium
30 bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof.

In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials. The term "pharmaceutical agent or drug" as used herein refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient. Other chemistry terms herein are used according to conventional usage in the art, as exemplified by *The McGraw-Hill Dictionary of Chemical Terms* (Parker, S., Ed., McGraw-Hill, San Francisco (1985)), incorporated herein by reference).

The term "antineoplastic agent" is used herein to refer to agents that have the functional property of inhibiting a development or progression of a neoplasm in a human, particularly a malignant (cancerous) lesion, such as a carcinoma, sarcoma, lymphoma, or leukemia. Inhibition of metastasis is frequently a property of antineoplastic agents.

The term patient includes human and veterinary subjects.

Human Anti-IGF-IR Antibodies and Characterization Thereof

Human antibodies avoid certain of the problems associated with antibodies that possess mouse or rat variable and/or constant regions. The presence of such mouse or rat derived sequences can lead to the rapid clearance of the antibodies or can lead to the generation of an immune response against the antibody by a patient. Therefore, in one embodiment, the invention provides humanized anti-IGF-IR antibodies. In a preferred embodiment, the invention provides fully human anti-IGF-IR antibodies by introducing human immunoglobulin genes into a rodent so that the rodent produces fully human antibodies. More preferred are fully human anti-human IGF-IR antibodies. Fully human anti-IGF-IR antibodies are expected to minimize the immunogenic and allergic responses intrinsic to mouse or mouse-derivatized monoclonal antibodies (Mabs) and thus to increase the efficacy and safety of the administered antibodies. The use of fully human antibodies can be expected to provide a substantial advantage in the treatment of chronic and recurring human diseases, such as inflammation and cancer, which may require repeated antibody

administrations. In another embodiment, the invention provides an anti-IGF-IR antibody that does not bind complement.

In a preferred embodiment, the anti-IGF-IR antibody is 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the anti-IGF-IR antibody comprises a light chain comprising an amino acid sequence selected from SEQ ID NO: 2, 6, 10, 14, 18 or 22, or one or more CDRs from these amino acid sequences. In another preferred embodiment, the anti-IGF-IR antibody comprises a heavy chain comprising an amino acid sequence selected from SEQ ID NO: 4, 8, 12, 16, 20 or 24 or one or more CDRs from these amino acid sequences.

10 *Class and Subclass of Anti-IGF-IR Antibodies*

The antibody may be an IgG, an IgM, an IgE, an IgA or an IgD molecule. In a preferred embodiment, the antibody is an IgG and is an IgG1, IgG2, IgG3 or IgG4 subtype. In a more preferred embodiment, the anti-IGF-IR antibody is subclass IgG2. In another preferred embodiment, the anti-IGF-IR antibody is the same class and subclass as antibody 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1, which is IgG2.

The class and subclass of anti-IGF-IR antibodies may be determined by any method known in the art. In general, the class and subclass of an antibody may be determined using antibodies that are specific for a particular class and subclass of antibody. Such antibodies are available commercially. The class and subclass can be determined by ELISA, Western Blot as well as other techniques. Alternatively, the class and subclass may be determined by sequencing all or a portion of the constant domains of the heavy and/or light chains of the antibodies, comparing their amino acid sequences to the known amino acid sequences of various class and subclasses of immunoglobulins, and determining the class and subclass of the antibodies.

Species and Molecule Selectivity

In another aspect of the invention, the anti-IGF-IR antibody demonstrates both species and molecule selectivity. In one embodiment, the anti-IGF-IR antibody binds to human, cynomolgous or rhesus IGF-IR. In a preferred embodiment, the anti-IGF-IR antibody does not bind to mouse, rat, guinea pig, dog or rabbit IGF-IR. In another preferred embodiment, the anti-IGF-IR antibody does not

bind to a New World monkey species such as a marmoset. Following the teachings of the specification, one may determine the species selectivity for the anti-IGF-IR antibody using methods well known in the art. For instance, one may determine species selectivity using Western blot, FACS, ELISA or RIA. In a preferred embodiment, one may determine the species selectivity using Western blot.

In another embodiment, the anti-IGF-IR antibody has a selectivity for IGF-IR that is at least 50 times greater than its selectivity for insulin receptor. In a preferred embodiment, the selectivity of the anti-IGF-IR antibody is more than 100 times greater than its selectivity for insulin receptor. In an even more preferred embodiment, the anti-IGF-IR antibody does not exhibit any appreciable specific binding to any other protein other than IGF-IR. One may determine the selectivity of the anti-IGF-IR antibody for IGF-IR using methods well known in the art following the teachings of the specification. For instance, one may determine the selectivity using Western blot, FACS, ELISA or RIA. In a preferred embodiment, one may determine the molecular selectivity using Western blot.

Binding Affinity of Anti-IGF-IR to IGF-IR

In another aspect of the invention, the anti-IGF-IR antibodies bind to IGF-IR with high affinity. In one embodiment, the anti-IGF-IR antibody binds to IGF-IR with a K_d of 1×10^{-8} M or less. In a more preferred embodiment, the antibody binds to IGF-IR with a K_d of 1×10^{-9} M or less. In an even more preferred embodiment, the antibody binds to IGF-IR with a K_d of 5×10^{-10} M or less. In another preferred embodiment, the antibody binds to IGF-IR with a K_d of 1×10^{-10} M or less. In another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_d as an antibody selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_d as an antibody that comprises one or more CDRs from an antibody selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In still another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_d as an antibody that comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24. In another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_d as an antibody that comprises one or more CDRs from an antibody that comprises one of

the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24.

In another aspect of the invention, the anti-IGF-IR antibody has a low dissociation rate. In one embodiment, the anti-IGF-IR antibody has an K_{off} of $1 \times 10^{-4} \text{ s}^{-1}$ or lower. In a preferred embodiment, the K_{off} is $5 \times 10^{-5} \text{ s}^{-1}$ or lower. In another preferred embodiment, the K_{off} is substantially the same as an antibody selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_{off} as an antibody that comprises one or more CDRs from an antibody selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In still another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_{off} as an antibody that comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24. In another preferred embodiment, the antibody binds to IGF-IR with substantially the same K_{off} as an antibody that comprises one or more CDRs from an antibody that comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22.

The binding affinity and dissociation rate of an anti-IGF-IR antibody to IGF-IR may be determined by any method known in the art. In one embodiment, the binding affinity can be measured by competitive ELISAs, RIAs or surface plasmon resonance, such as BIAcore. The dissociation rate can also be measured by surface plasmon resonance. In a more preferred embodiment, the binding affinity and dissociation rate is measured by surface plasmon resonance. In an even more preferred embodiment, the binding affinity and dissociation rate is measured using a BIAcore. An example of determining binding affinity and dissociation rate is described below in Example II.

Half-Life of Anti-IGF-IR Antibodies

According to another object of the invention, the anti-IGF-IR antibody has a half-life of at least one day *in vitro* or *in vivo*. In a preferred embodiment, the antibody or portion thereof has a half-life of at least three days. In a more preferred embodiment, the antibody or portion thereof has a half-life of four days or longer. In another embodiment, the antibody or portion thereof has a half-life of eight days or longer. In another embodiment, the antibody or antigen-binding portion thereof is derivatized or modified such that it has a longer half-life, as discussed below. In

another preferred embodiment, the antibody may contain point mutations to increase serum half life, such as described WO 00/09560, published February 24, 2000.

The antibody half-life may be measured by any means known to one having ordinary skill in the art. For instance, the antibody half life may be measured by Western blot, ELISA or RIA over an appropriate period of time. The antibody half-life may be measured in any appropriate animals, e.g., a monkey, such as a cynomologous monkey, a primate or a human.

*Identification of IGF-IR Epitopes
Recognized by Anti-IGF-IR Antibody*

The invention also provides an anti-IGF-IR antibody that binds the same antigen or epitope as a human anti-IGF-IR antibody. Further, the invention provides an anti-IGF-IR antibody that cross-competes with a human anti-IGF-IR antibody. In a preferred embodiment, the human anti-IGF-IR antibody is 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the human anti-IGF-IR comprises one or more CDRs from an antibody selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In still another preferred embodiment, the human anti-IGF-IR comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24. In another preferred embodiment, the human anti-IGF-IR comprises one or more CDRs from an antibody that comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24. In a highly preferred embodiment, the anti-IGF-IR antibody is another human antibody.

One may determine whether an anti-IGF-IR antibody binds to the same antigen using a variety of methods known in the art. For instance, one may determine whether a test anti-IGF-IR antibody binds to the same antigen by using an anti-IGF-IR antibody to capture an antigen that is known to bind to the anti-IGF-IR antibody, such as IGF-IR, eluting the antigen from the antibody, and then determining whether the test antibody will bind to the eluted antigen. One may determine whether an antibody binds to the same epitope as an anti-IGF-IR antibody by binding the anti-IGF-IR antibody to IGF-IR under saturating conditions, and then measuring the ability of the test antibody to bind to IGF-IR. If the test antibody is able to bind to the IGF-IR at the same time as the anti-IGF-IR antibody, then the test antibody binds to a different epitope as the anti-IGF-IR antibody. However, if the test antibody is not

able to bind to the IGF-IR at the same time, then the test antibody binds to the same, epitope as the human anti-IGF-IR antibody. This experiment may be performed using ELISA, RIA or surface plasmon resonance. In a preferred embodiment, the experiment is performed using surface plasmon resonance. In a more preferred
5 embodiment, BIAcore is used. One may also determine whether an anti-IGF-IR antibody cross-competes with an anti-IGF-IR antibody. In a preferred embodiment, one may determine whether an anti-IGF-IR antibody cross-competes with another by using the same method that is used to measure whether the anti-IGF-IR antibody is able to bind to the same epitope as another anti-IGF-IR antibody.

10 *Light and Heavy Chain Usage*

The invention also provides an anti-IGF-IR antibody that comprises variable sequences encoded by a human κ gene. In a preferred embodiment, the variable sequences are encoded by either the V κ A27, A30 or O12 gene family. In a preferred embodiment, the variable sequences are encoded by a human V κ A30 gene
15 family. In a more preferred embodiment, the light chain comprises no more than ten amino acid substitutions from the germline V κ A27, A30 or O12, preferably no more than six amino acid substitutions, and more preferably no more than three amino acid substitutions. In a preferred embodiment, the amino acid substitutions are conservative substitutions.

20 SEQ ID NOS: 2, 6, 10, 14, 18 and 22 provide the amino acid sequences of the variable regions of six anti-IGF-IR κ light chains. SEQ ID NOS: 38, 40 and 42 provide the amino acid sequences of the three germline κ light chains from which the six anti-IGF-IR κ light chains are derived. Figs. 1A-1C show the alignments of the nucleotide sequences of the variable regions of the light chains of
25 the six anti-IGF-IR antibodies to each other and to the germline sequences from which they are derived. Following the teachings of this specification, one of ordinary skill in the art could determine the encoded amino acid sequence of the six anti-IGF-IR κ light chains and the germline κ light chains and determine the differences between the germline sequences and the antibody sequences.

30 In a preferred embodiment, the VL of the anti-IGF-IR antibody contains the same amino acid substitutions, relative to the germline amino acid sequence, as any one or more of the VL of antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. For example, the VL of the anti-IGF-IR antibody may contain

one or more amino acid substitutions that are the same as those present in antibody 2.13.2, another amino acid substitution that is the same as that present in antibody 2.14.3, and another amino acid substitution that is the same as antibody 4.9.2. In this manner, one can mix and match different features of antibody binding in order to alter, e.g., the affinity of the antibody for IGF-IR or its dissociation rate from the antigen. In another embodiment, the amino acid substitutions are made in the same position as those found in any one or more of the VL of antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1, but conservative amino acid substitutions are made rather than using the same amino acid. For example, if the amino acid substitution compared to the germline in one of the antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1 is glutamate, one may conservatively substitute aspartate. Similarly, if the amino acid substitution is serine, one may conservatively substitute threonine.

In another preferred embodiment, the light chain comprises an amino acid sequence that is the same as the amino acid sequence of the VL of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another highly preferred embodiment, the light chain comprises amino acid sequences that are the same as the CDR regions of the light chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the light chain comprises an amino acid sequence from at least one CDR region of the light chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the light chain comprises amino acid sequences from CDRs from different light chains. In a more preferred embodiment, the CDRs from different light chains are obtained from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the light chain comprises an amino acid sequence selected from SEQ ID NOS: 2, 6, 10, 14, 18 or 22. In another embodiment, the light chain comprises an amino acid sequence encoded by a nucleic acid sequence selected from SEQ ID NOS: 1, 5, 9, 13, 17 or 21, or a nucleic acid sequence that encodes an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions therefrom. Preferably, the amino acid substitutions are conservative amino acid substitutions. In another embodiment, the antibody or portion thereof comprises a lambda light chain.

The present invention also provides an anti-IGF-IR antibody or portion thereof comprises a human heavy chain or a sequence derived from a human heavy

chain. In one embodiment, the heavy chain amino acid sequence is derived from a human V_H DP-35, DP-47, DP-70, DP-71 or VIV-4/4.35 gene family. In a preferred embodiment, the heavy chain amino acid sequence is derived from a human V_H DP-47 gene family. In a more preferred embodiment, the heavy chain comprises no more than eight amino acid changes from germline V_H DP-35, DP-47, DP-70, DP-71 or VIV-4/4.35, more preferably no more than six amino acid changes, and even more preferably no more than three amino acid changes.

SEQ ID NOS: 4, 8, 12, 16, 20 and 24 provide the amino acid sequences of the variable regions of six anti-IGF-IR heavy chains. SEQ ID NOS: 30, 32, 34, 36 and 44 provide the amino acid sequences and SEQ ID NOS: 29, 31, 33, 35 and 43 provide the nucleotide sequences of the germline heavy chains DP-35, DP-47, DP-70, DP-71 and VIV-4, respectively. Figs. 2A-2D show the alignments of the amino acid sequences of the variable region of the six anti-IGF-IR antibodies to their corresponding germline sequences. Following the teachings of this specification, one of ordinary skill in the art could determine the encoded amino acid sequence of the six anti-IGF-IR heavy chains and the germline heavy chains and determine the differences between the germline sequences and the antibody sequences.

In a preferred embodiment, the V_H of the anti-IGF-IR antibody contains the same amino acid substitutions, relative to the germline amino acid sequence, as any one or more of the V_H of antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. Similar to what was discussed above, the V_H of the anti-IGF-IR antibody may contain one or more amino acid substitutions that are the same as those present in antibody 2.13.2, another amino acid substitution that is the same as that present in antibody 2.14.3, and another amino acid substitution that is the same as antibody 4.9.2. In this manner, one can mix and match different features of antibody binding in order to alter, e.g., the affinity of the antibody for IGF-IR or its dissociation rate from the antigen. In another embodiment, the amino acid substitutions are made in the same position as those found in any one or more of the V_H of antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.17.3, 4.9.2 or 6.1.1, but conservative amino acid substitutions are made rather than using the same amino acid.

In another preferred embodiment, the heavy chain comprises an amino acid sequence that is the same as the amino acid sequence of the V_H of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another highly preferred embodiment, the

heavy chain comprises amino acid sequences that are the same as the CDR regions of the heavy chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the heavy chain comprises an amino acid sequence from at least one CDR region of the heavy chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the heavy chain comprises amino acid sequences from CDRs from different heavy chains. In a more preferred embodiment, the CDRs from different heavy chains are obtained from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred embodiment, the heavy chain comprises an amino acid sequence selected from SEQ ID NOS: 4, 8, 12, 16, 20 or 24. In another embodiment, the heavy chain comprises an amino acid sequence encoded by a nucleic acid sequence selected from SEQ ID NOS: 3, 7, 11, 15, 19 or 23, or a nucleic acid sequence that encodes an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions therefrom. In another embodiment, the substitutions are conservative amino acid substitutions.

15 Inhibition of IGF-IR Activity by Anti-IGF-IR Antibody

Inhibition of IGF-I Binding to IGF-IR

In another embodiment, the invention provides an anti-IGF-IR antibody that inhibits the binding of IGF-I to IGF-IR or the binding of IGF-II to IGF-IR. In a preferred embodiment, the IGF-IR is human. In another preferred embodiment, the anti-IGF-IR antibody is a human antibody. In another embodiment, the antibody or portion thereof inhibits binding between IGF-IR and IGF-I with an IC_{50} of no more than 100 nM. In a preferred embodiment, the IC_{50} is no more than 10 nM. In a more preferred embodiment, the IC_{50} is no more than 5 nM. The IC_{50} can be measured by any method known in the art. Typically, an IC_{50} can be measured by ELISA or RIA. In a preferred embodiment, the IC_{50} is measured by RIA.

In another embodiment, the invention provides an anti-IGF-IR antibody that prevents activation of the IGF-IR in the presence of IGF-I. In a preferred embodiment, the anti-IGF-IR antibody inhibits IGF-IR-induced tyrosine phosphorylation that occurs upon occupancy of the receptor. In another preferred embodiment, the anti-IGF-IR antibody inhibits downstream cellular events from occurring. For instance, the anti-IGF-IR can inhibit tyrosine phosphorylation of Shc and insulin receptor substrate (IRS) 1 and 2, all of which are normally phosphorylated

when cells are treated with IGF-I (Kim et al., J. Biol. Chem. 273: 34543-34550, 1998). One can determine whether an anti-IGF-IR antibody can prevent activation of IGF-IR in the presence of IGF-1 by determining the levels of autophosphorylation for IGF-IR, Shc, IRS-1 or IRS-2 by Western blot or immunoprecipitation. In a preferred
5 embodiment, one would determine the levels of autophosphorylation of IGF-IR by Western blot. *See, e.g.*, Example VII.

In another aspect of the invention, the antibody causes the downregulation of IGF-IR from a cell treated with the antibody. In one embodiment, the IGF-IR is internalized into the cytoplasm of the cell. After the anti-IGF-IR
10 antibody binds to IGF-IR, the antibody is internalized, as shown by confocal microscopy. Without wishing to be bound to any theory, it is believed that the antibody-IGF-IR complex is internalized into a lysosome and degraded. One may measure the downregulation of IGF-IR by any method known in the art including immunoprecipitation, confocal microscopy or Western blot. *See, e.g.*, Example VII.
15 In a preferred embodiment, the antibody is selected 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding region thereof.

Activation of IGF-IR by Anti-IGF-IR Antibody

Another aspect of the present invention involves activating anti-IGF-IR antibodies. An activating antibody differs from an inhibiting antibody because it
20 amplifies or substitutes for the effects of IGF-I on IGF-IR. In one embodiment, the activating antibody is able to bind to IGF-IR and cause it to be activated in the absence of IGF-I. This type of activating antibody is essentially a mimic of IGF-I. In another embodiment, the activating antibody amplifies the effect of IGF-I on IGF-IR. This type of antibody does not activate IGF-IR by itself, but rather increases the
25 activation of IGF-IR in the presence of IGF-I. A mimic anti-IGF-IR antibody may be easily distinguished from an amplifying anti-IGF-IR antibody by treating cells *in vitro* with an antibody in the presence or absence of low levels of IGF-I. If the antibody is able to cause IGF-IR activation in the absence of IGF-I, e.g., it increases IGF-IR tyrosine phosphorylation, then the antibody is a mimic antibody. If the antibody
30 cannot cause IGF-IR activation in the absence of IGF-I but is able to amplify the amount of IGF-IR activation, then the antibody is an amplifying antibody. In a preferred embodiment, the activating antibody is 4.17.3. In another preferred embodiment, the antibody comprises one or more CDRs from 4.17.3. In another

preferred embodiment, the antibody is derived from either or both of the germline sequences O12 (light chain) and/or D71 (heavy chain).

Inhibition of IGF-IR Tyrosine Phosphorylation, IGF-IR Levels and Tumor Cell Growth *In Vivo* by Anti-IGF-IR Antibodies

5 Another embodiment of the invention provides an anti-IGF-IR antibody that inhibits IGF-IR tyrosine phosphorylation and receptor levels *in vivo*. In one embodiment, administration of anti-IGF-IR antibody to an animal causes a reduction in IGF-IR phosphotyrosine signal in IGF-IR-expressing tumors. In a preferred embodiment, the anti-IGF-IR antibody causes a reduction in
10 phosphotyrosine signal by at least 20%. In a more preferred embodiment, the anti-IGF-IR antibody causes a decrease in phosphotyrosine signal by at least 60%, more preferably 50%. In an even more preferred embodiment, the antibody causes a decrease in phosphotyrosine signal of at least 40%, more preferably 30%, even more preferably 20%. In a preferred embodiment, the antibody is administered
15 approximately 24 hours before the levels of tyrosine phosphorylation are measured. The levels of tyrosine phosphorylation may be measured by any method known in the art, such as those described *infra*. See, e.g., Example III and Figure 5. In a preferred embodiment, the antibody is selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding portion thereof.

20 In another embodiment, administration of anti-IGF-IR antibody to an animal causes a reduction in IGF-IR levels in IGF-IR-expressing tumors. In a preferred embodiment, the anti-IGF-IR antibody causes a reduction in receptor levels by at least 20% compared to an untreated animal. In a more preferred embodiment, the anti-IGF-IR antibody causes a decrease in receptor levels to at least 60%, more
25 preferably 50% of the receptor levels in an untreated animal. In an even more preferred embodiment, the antibody causes a decrease in receptor levels to at least 40%, more preferably 30%. In a preferred embodiment, the antibody is administered approximately 24 hours before the IGF-IR levels are measured. The IGF-IR levels may be measured by any method known in the art, such as those described *infra*. See,
30 e.g., Example VIII and Figure 6. In a preferred embodiment, the antibody is selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding portion thereof.

In another embodiment, an anti-IGF-IR antibody inhibits tumor cell growth *in vivo*. The tumor cell may be derived from any cell type including, without limitation, epidermal, epithelial, endothelial, leukemia, sarcoma, multiple myeloma or mesodermal cells. Examples of tumor cells include A549 (non-small cell lung carcinoma) cells, MCF-7 cells, Colo 205 cells, 3T3/IGF-IR cells and A431 cells. In a preferred embodiment, the antibody inhibits tumor cell growth as compared to the growth of the tumor in an untreated animal. In a more preferred embodiment, the antibody inhibits tumor cell growth by 50%. In an even more preferred embodiment, the antibody inhibits tumor cell growth by 60%, 65%, 70% or 75%. In one embodiment, the inhibition of tumor cell growth is measured at least 7 days after the animals have started treatment with the antibody. In a more preferred embodiment, the inhibition of tumor cell growth is measured at least 14 days after the animals have started treatment with the antibody. In another preferred embodiment, another antineoplastic agent is administered to the animal with the anti-IGF-IR antibody. In a preferred embodiment, the antineoplastic agent is able to further inhibit tumor cell growth. In an even more preferred embodiment, the antineoplastic agent is adriamycin, taxol, tamoxifen, 5-fluorodeoxyuridine (5-FU) or CP-358,774. In a preferred embodiment, the co-administration of an antineoplastic agent and the anti-IGF-IR antibody inhibits tumor cell growth by at least 50%, more preferably 60%, 65%, 70% or 75%, more preferably 80%, 85% or 90% after a period of 22-24 days. See, e.g., Fig. 7 and Example IX. In a preferred embodiment, the antibody is selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding portion thereof.

Induction of Apoptosis by Anti-IGF-IR Antibodies

Another aspect of the invention provides an anti-IGF-IR antibody that induces cell death. In one embodiment, the antibody causes apoptosis. The antibody may induce apoptosis either *in vivo* or *in vitro*. In general, tumor cells are more sensitive to apoptosis than normal cells, such that administration of an anti-IGF-IR antibody causes apoptosis of a tumor cell preferentially to that of a normal cell. In another embodiment, the administration of an anti-IGF-IR antibody decreases levels of an enzyme, *akt*, which is involved in the phosphatidyl inositol (PI) kinase pathway. The PI kinase pathway, in turn, is involved in the cell proliferation and prevention of apoptosis. Thus, inhibition of *akt* can cause apoptosis. In a more preferred

embodiment, the antibody is administered *in vivo* to cause apoptosis of an IGF-IR-expressing cell. In a preferred embodiment, the antibody is selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding portion thereof.

5 Methods of Producing Antibodies and Antibody-Producing Cell Lines

Immunization

In one embodiment of the instant invention, human antibodies are produced by immunizing a non-human animal comprising some or all of the human immunoglobulin locus with an IGF-IR antigen. In a preferred embodiment, the non-
10 human animal is a XENOMOUSE™, which is an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production. See, e.g., Green et al. *Nature Genetics* 7:13-21 (1994) and United States Patents 5,916,771, 5,939,598, 5,985,615, 5,998,209, 6,075,181, 6,091,001, 6,114,598 and 6,130,364. See also WO 91/10741, published July 25,
15 1991, WO 94/02602, published February 3, 1994, WO 96/34096 and WO 96/33735, both published October 31, 1996, WO 98/16654, published April 23, 1998, WO 98/24893, published June 11, 1998, WO 98/50433, published November 12, 1998, WO 99/45031, published September 10, 1999, WO 99/53049, published October 21, 1999, WO 00 09560, published February 24, 2000 and WO 00/037504, published
20 June 29, 2000. The XENOMOUSE™ produces an adult-like human repertoire of fully human antibodies, and generates antigen-specific human Mabs. A second generation XENOMOUSE™ contains approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and κ light chain loci. See Mendez et al.
25 *Nature Genetics* 15:146-156 (1997), Green and Jakobovits *J. Exp. Med.* 188:483-495 (1998), the disclosures of which are hereby incorporated by reference.

The invention also provides a method for making anti-IGF-IR antibodies from non-human, non-mouse animals by immunizing non-human
transgenic animals that comprise human immunoglobulin loci. One may produce
30 such animals using the methods described immediately above. The methods disclosed in these patents may modified as described in United States Patent 5,994,619. In a

preferred embodiment, the non-human animals may be rats, sheep, pigs, goats, cattle or horses.

In another embodiment, the non-human animal comprising human immunoglobulin gene loci are animals that have a "minilocus" of human immunoglobulins. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of individual genes from the Ig locus. Thus, one or more V_H genes, one or more D_H genes, one or more J_H genes, a mu constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This approach is described, *inter alia*, in U.S. Patent No. 5,545,807, 5,545,806, 5,625,825, 5,625,126, 5,633,425, 5,661,016, 5,770,429, 5,789,650, 5,814,318, 5,591,669, 5,612,205, 5,721,367, 5,789,215, and 5,643,763, hereby incorporated by reference.

An advantage of the minilocus approach is the rapidity with which constructs including portions of the Ig locus can be generated and introduced into animals. However, a potential disadvantage of the minilocus approach is that there may not be sufficient immunoglobulin diversity to support full B-cell development, such that there may be lower antibody production.

In order to produce a human anti-IGF-IR antibody, a non-human animal comprising some or all of the human immunoglobulin loci is immunized with an IGF-IR antigen and the antibody or the antibody-producing cell is isolated from the animal. The IGF-IR antigen may be isolated and/or purified IGF-IR and is preferably a human IGF-IR. In another embodiment, the IGF-IR antigen is a fragment of IGF-IR, preferably the extracellular domain of IGF-IR. In another embodiment, the IGF-IR antigen is a fragment that comprises at least one epitope of IGF-IR. In another embodiment, the IGF-IR antigen is a cell that expresses IGF-IR on its cell surface, preferably a cell that overexpresses IGF-IR on its cell surface.

Immunization of animals may be done by any method known in the art. See, e.g., Harlow and Lane, Antibodies: A Laboratory Manual, New York: Cold Spring Harbor Press, 1990. Methods for immunizing non-human animals such as mice, rats, sheep, goats, pigs, cattle and horses are well known in the art. See, e.g., Harlow and Lane and United States Patent 5,994,619. In a preferred embodiment, the IGF-IR antigen is administered with an adjuvant to stimulate the immune response. Such adjuvants include complete or incomplete Freund's adjuvant, RIBI (muramyl

dipeptides) or ISCOM (immunostimulating complexes). Such adjuvants may protect the polypeptide from rapid dispersal by sequestering it in a local deposit, or they may contain substances that stimulate the host to secrete factors that are chemotactic for macrophages and other components of the immune system. Preferably, if a polypeptide is being administered, the immunization schedule will involve two or more administrations of the polypeptide, spread out over several weeks.

Example I provides an protocol for immunizing a XENOMOUSE[™] with full-length human IGF-IR in phosphate-buffered saline.

Production of Antibodies and Antibody-Producing Cell Lines

After immunization of an animal with an IGF-IR antigen, antibodies and/or antibody-producing cells may be obtained from the animal. An anti-IGF-IR antibody-containing serum is obtained from the animal by bleeding or sacrificing the animal. The serum may be used as it is obtained from the animal, an immunoglobulin fraction may be obtained from the serum, or the anti-IGF-IR antibodies may be purified from the serum. Serum or immunoglobulins obtained in this manner are polyclonal, which are disadvantageous because the amount of antibodies that can be obtained is limited and the polyclonal antibody has a heterogeneous array of properties.

In another embodiment, antibody-producing immortalized hybridomas may be prepared from the immunized animal. After immunization, the animal is sacrificed and the splenic B cells are fused to immortalized myeloma cells as is well-known in the art. See, e.g., Harlow and Lane, *supra*. In a preferred embodiment, the myeloma cells do not secrete immunoglobulin polypeptides (a non-secretory cell line). After fusion and antibiotic selection, the hybridomas are screened using IGF-IR, a portion thereof, or a cell expressing IGF-IR. In a preferred embodiment, the initial screening is performed using an enzyme-linked immunoassay (ELISA) or a radioimmunoassay (RIA), preferably an ELISA. An example of ELISA screening is provided in WO 00/37504, herein incorporated by reference.

In another embodiment, antibody-producing cells may be prepared from a human who has an autoimmune disorder and who expresses anti-IGF-IR antibodies. Cells expressing the anti-IGF-IR antibodies may be isolated by isolating white blood cells and subjecting them to fluorescence-activated cell sorting (FACS) or by panning on plates coated with IGF-IR or a portion thereof. These cells may be

fused with a human non-secretory myeloma to produce human hybridomas expressing human anti-IGF-IR antibodies. In general, this is a less preferred embodiment because it is likely that the anti-IGF-IR antibodies will have a low affinity for IGF-IR.

Anti-IGF-IR antibody-producing hybridomas are selected, cloned and
5 further screened for desirable characteristics, including robust hybridoma growth, high antibody production and desirable antibody characteristics, as discussed further below. Hybridomas may be cultured and expanded *in vivo* in syngeneic animals, in animals that lack an immune system, e.g., nude mice, or in cell culture *in vitro*. Methods of selecting, cloning and expanding hybridomas are well known to those of
10 ordinary skill in the art.

Preferably, the immunized animal is a non-human animal that expresses human immunoglobulin genes and the splenic B cells are fused to a myeloma derived from the same species as the non-human animal. More preferably, the immunized animal is a XENOMOUSE™ and the myeloma cell line is a non-
15 secretory mouse myeloma, such as the myeloma cell line is NSO-bcl2. See, e.g., Example I.

In one aspect, the invention provides hybridomas are produced that produce human anti-IGF-IR antibodies. In a preferred embodiment, the hybridomas are mouse hybridomas, as described above. In another preferred embodiment, the
20 hybridomas are produced in a non-human, non-mouse species such as rats, sheep, pigs, goats, cattle or horses. In another embodiment, the hybridomas are human hybridomas, in which a human non-secretory myeloma is fused with a human cell expressing an anti-IGF-IR antibody.

25 Nucleic Acids, Vectors, Host Cells and Recombinant Methods of Making Antibodies

Nucleic Acids

Nucleic acid molecules encoding anti-IGF-IR antibodies of the invention are provided. In one embodiment, the nucleic acid molecule encodes a heavy and/or light chain of an anti-IGF-IR immunoglobulin. In a preferred
30 embodiment, a single nucleic acid molecule encodes a heavy chain of an anti-IGF-IR immunoglobulin and another nucleic acid molecule encodes the light chain of an anti-IGF-IR immunoglobulin. In a more preferred embodiment, the encoded

immunoglobulin is a human immunoglobulin, preferably a human IgG. The encoded light chain may be a λ chain or a κ chain, preferably a κ chain.

The nucleic acid molecule encoding the variable region of the light chain may be derived from the A30, A27 or O12 V κ gene. In a preferred
5 embodiment, the light chain is derived from the A30 V κ gene. In another preferred embodiment, the nucleic acid molecule encoding the light chain comprises the joining region derived from J κ 1, J κ 2 or J κ 4. In an even more preferred embodiment, the nucleic acid molecule encoding the light chain contains no more than ten amino acid changes from the germline A30 V κ gene, preferably no more than six amino acid
10 changes, and even more preferably no more than three amino acid changes.

The invention provides a nucleic acid molecule that encodes a variable region of the light chain (VL) containing at least three amino acid changes compared to the germline sequence, wherein the amino acid changes are identical to the amino acid changes from the germline sequence from the VL of one of the antibodies 2.12.1,
15 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. The invention also provides a nucleic acid molecule comprising a nucleic acid sequence that encodes the amino acid sequence of the variable region of the light chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. The invention also provides a nucleic acid molecule comprising a nucleic acid sequence that encodes the amino acid sequence of one or more of the CDRs of any
20 one of the light chains of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In a preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of all of the CDRs of any one of the light chains of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid
25 sequence of one of SEQ ID NOS: 2, 6, 10, 14, 18 or 22 or comprises a nucleic acid sequence of one of SEQ ID NOS: 1, 5, 9, 13, 17 or 21. In another preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of one or more of the CDRs of any one of SEQ ID NOS: 2, 6, 10, 14, 18 or 22 or comprises a nucleic acid sequence of one or more of
30 the CDRs of any one of SEQ ID NOS: 1, 5, 9, 13, 17 or 21. In a more preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of all of the CDRs of any one of SEQ ID NOS: 2, 6,

10, 14, 18 or 22 or comprises a nucleic acid sequence of all the CDRs of any one of SEQ ID NOS: 1, 5, 9, 13, 17 or 21.

The invention also provides a nucleic acid molecules that encodes an amino acid sequence of a VL that has an amino acid sequence that is at least 70%,
5 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a VL described above, particularly to a VL that comprises an amino acid sequence of one of SEQ ID NOS: 2, 6, 10, 14, 18 or 22. The invention also provides a nucleic acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence of one of SEQ ID NOS: 1, 5, 9, 13, 17 or 21. In another
10 embodiment, the invention provides a nucleic acid molecule encoding a VL that hybridizes under highly stringent conditions to a nucleic acid molecule encoding a VL as described above, particularly a nucleic acid molecule that comprises a nucleic acid sequence encoding an amino acid sequence of SEQ ID NOS: 2, 6, 10, 14, 18 or 22. The invention also provides a nucleic acid sequence encoding an VL that hybridizes
15 under highly stringent conditions to a nucleic acid molecule comprising a nucleic acid sequence of one of SEQ ID NOS: 1, 5, 9, 13, 17 or 21.

The invention also provides a nucleic acid molecule encoding the variable region of the heavy chain (VH) is derived from the DP-35, DP-47, DP-71 or VIV-4/4.35 VH gene, preferably the DP-35 VH gene. In another preferred
20 embodiment, the nucleic acid molecule encoding the VH comprises the joining region derived from JH6 or JH5, more preferably JH6. In another preferred embodiment, the D segment is derived from 3-3, 6-19 or 4-17. In an even more preferred embodiment, the nucleic acid molecule encoding the VH contains no more than ten amino acid changes from the germline DP-47 gene, preferably no more than six amino acid
25 changes, and even more preferably no more than three amino acid changes. In a highly preferred embodiment, the nucleic acid molecule encoding the VH contains at least one amino acid change compared to the germline sequence, wherein the amino acid change is identical to the amino acid change from the germline sequence from the heavy chain of one of the antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or
30 6.1.1. In an even more preferred embodiment, the VH contains at least three amino acid changes compared to the germline sequences, wherein the changes are identical to those changes from the germline sequence from the VH of one of the antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1.

In one embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of the VH of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of one or
5 more of the CDRs of the heavy chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In a preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequences of all of the CDRs of the heavy chain of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another preferred
10 embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of one of SEQ ID NOS: 4, 8, 12, 16, 20 or 24 or that comprises a nucleic acid sequence of one of SEQ ID NOS: 3, 7, 11, 15, 19 or 23. In another preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequence of one or more of the CDRs of any
15 one of SEQ ID NOS: 4, 8, 12, 16, 20 or 24 or comprises a nucleic acid sequence of one or more of the CDRs of any one of SEQ ID NOS: 3, 7, 11, 15, 19 or 23. In a preferred embodiment, the nucleic acid molecule comprises a nucleic acid sequence that encodes the amino acid sequences of all of the CDRs of any one of SEQ ID NOS: 4, 8, 12, 16, 20 or 24 or comprises a nucleic acid sequence of all of the CDRs of any
20 one of SEQ ID NOS: 3, 7, 11, 15, 19 or 23.

In another embodiment, the nucleic acid molecule encodes an amino acid sequence of a VH that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to one of the amino acid sequences encoding a VH as described immediately above, particularly to a VH that comprises an amino acid sequence of one of SEQ ID NOS: 4, 8, 12, 16, 20 or 24. The invention also provides a nucleic
25 acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence of one of SEQ ID NOS: 3, 7, 11, 15, 19 or 23. In another embodiment, the nucleic acid molecule encoding a VH is one that hybridizes under highly stringent conditions to a nucleic acid sequence encoding a
30 VH as described above, particularly to a VH that comprises an amino acid sequence of one of SEQ ID NOS: 4, 8, 12, 16, 20 or 24. The invention also provides a nucleic acid sequence encoding a VH that hybridizes under highly stringent conditions to a nucleic acid molecule comprising a nucleic acid sequence of one of SEQ ID NOS: 3, 7, 11, 15, 19 or 23.

The nucleic acid molecule encoding either or both of the entire heavy and light chains of an anti-IGF-IR antibody or the variable regions thereof may be obtained from any source that produces an anti-IGF-IR antibody. Methods of isolating mRNA encoding an antibody are well-known in the art. See, e.g., Sambrook et al. The mRNA may be used to produce cDNA for use in the polymerase chain reaction (PCR) or cDNA cloning of antibody genes. In one embodiment of the invention, the nucleic acid molecules may be obtained from a hybridoma that expresses an anti-IGF-IR antibody, as described above, preferably a hybridoma that has as one of its fusion partners a transgenic animal cell that expresses human immunoglobulin genes, such as a XENOMOUSE™, non-human mouse transgenic animal or a non-human, non-mouse transgenic animal. In another embodiment, the hybridoma is derived from a non-human, non-transgenic animal, which may be used, e.g., for humanized antibodies.

A nucleic acid molecule encoding the entire heavy chain of an anti-IGF-IR antibody may be constructed by fusing a nucleic acid molecule encoding the variable domain of a heavy chain or an antigen-binding domain thereof with a constant domain of a heavy chain. Similarly, a nucleic acid molecule encoding the light chain of an anti-IGF-IR antibody may be constructed by fusing a nucleic acid molecule encoding the variable domain of a light chain or an antigen-binding domain thereof with a constant domain of a light chain. The nucleic acid molecules encoding the VH and VL chain may be converted to full-length antibody genes by inserting them into expression vectors already encoding heavy chain constant and light chain constant regions, respectively, such that the VH segment is operatively linked to the heavy chain constant region (CH) segment(s) within the vector and the VL segment is operatively linked to the light chain constant region (CL) segment within the vector. Alternatively, the nucleic acid molecules encoding the VH or VL chains are converted into full-length antibody genes by linking, e.g., ligating, the nucleic acid molecule encoding a VH chain to a nucleic acid molecule encoding a CH chain using standard molecular biological techniques. The same may be achieved using nucleic acid molecules encoding VL and CL chains. The sequences of human heavy and light chain constant region genes are known in the art. See, e.g., Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed., NIH Publ. No. 91-3242, 1991. Nucleic acid molecules encoding the full-length heavy and/or light chains may then be

expressed from a cell into which they have been introduced and the anti-IGF-IR antibody isolated.

In a preferred embodiment, the nucleic acid encoding the variable region of the heavy chain encodes the amino acid sequence of SEQ ID NOS: 4, 8, 12, 16, 20 or 24, and the nucleic acid molecule encoding the variable region of the light chains encodes the amino acid sequence of SEQ ID NOS: 2, 6, 10, 14, 18 or 22. SEQ ID NO: 28 depicts the amino acid sequence and SEQ ID NO: 27 depicts the nucleic acid sequence encoding the constant region of the heavy chain of the anti-IGF-IR antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 and 6.1.1. SEQ ID NO: 26 depicts the amino acid sequence and SEQ ID NO: 25 depicts the nucleic acid sequence encoding the constant region of the light chain of the anti-IGF-IR antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 and 6.1.1. Thus, in a preferred embodiment, the nucleic acid molecule encoding the constant domain of the heavy chain encodes SEQ ID NO: 28, and the nucleic acid molecule encoding the constant domain of the light chain encodes SEQ ID NO: 26. In a more preferred embodiment, the nucleic acid molecule encoding the constant domain of the heavy chain has the nucleic acid sequence of SEQ ID NO: 27, and the nucleic acid molecule encoding the constant domain has the nucleic acid sequence of SEQ ID NO: 25.

In another embodiment, a nucleic acid molecule encoding either the heavy chain of an anti-IGF-IR antibody or an antigen-binding domain thereof, or the light chain of an anti-IGF-IR antibody or an antigen-binding domain thereof may be isolated from a non-human, non-mouse animal that expresses human immunoglobulin genes and has been immunized with an IGF-IR antigen. In other embodiment, the nucleic acid molecule may be isolated from an anti-IGF-IR antibody-producing cell derived from a non-transgenic animal or from a human patient who produces anti-IGF-IR antibodies. Methods of isolating mRNA from the anti-IGF-IR antibody-producing cells may be isolated by standard techniques, cloned and/or amplified using PCR and library construction techniques, and screened using standard protocols to obtain nucleic acid molecules encoding anti-IGF-IR heavy and light chains.

The nucleic acid molecules may be used to recombinantly express large quantities of anti-IGF-IR antibodies, as described below. The nucleic acid molecules may also be used to produce chimeric antibodies, single chain antibodies, immunoadhesins, diabodies, mutated antibodies and antibody derivatives, as

described further below. If the nucleic acid molecules are derived from a non-human, non-transgenic animal, the nucleic acid molecules may be used for antibody humanization, also as described below.

In another embodiment, the nucleic acid molecules of the invention
5 may be used as probes or PCR primers for specific antibody sequences. For instance, a nucleic acid molecule probe may be used in diagnostic methods or a nucleic acid molecule PCR primer may be used to amplify regions of DNA that could be used, *inter alia*, to isolate nucleic acid sequences for use in producing variable domains of anti-IGF-IR antibodies. In a preferred embodiment, the nucleic acid molecules are
10 oligonucleotides. In a more preferred embodiment, the oligonucleotides are from highly variable regions of the heavy and light chains of the antibody of interest. In an even more preferred embodiment, the oligonucleotides encode all or a part of one or more of the CDRs.

Vectors

15 The invention provides vectors comprising the nucleic acid molecules of the invention that encode the heavy chain or the antigen-binding portion thereof. The invention also provides vectors comprising the nucleic acid molecules of the invention that encode the light chain or antigen-binding portion thereof. The invention also provides vectors comprising nucleic acid molecules encoding fusion
20 proteins, modified antibodies, antibody fragments, and probes thereof.

To express the antibodies, or antibody portions of the invention, DNAs encoding partial or full-length light and heavy chains, obtained as described above, are inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. Expression vectors include
25 plasmids, retroviruses, cosmids, YACs, EBV derived episomes, and the like. The antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host
30 cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector. In a preferred embodiment, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the

antibody gene fragment and vector, or blunt end ligation if no restriction sites are present).

A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed, as described above. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The recombinant expression vector can also encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene may be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

In addition to the antibody chain genes, the recombinant expression vectors of the invention carry regulatory sequences that control the expression of the antibody chain genes in a host cell. It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from retroviral LTRs, cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)), polyoma and strong mammalian promoters such as native immunoglobulin and actin promoters. For further description of viral regulatory elements, and sequences thereof, see e.g., U.S. Pat. No. 5,168,062 by Stinski, U.S. Pat. No. 4,510,245 by Bell et al. and U.S. Pat. No. 4,968,615 by Schaffner et al.

In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of

replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see e.g., U.S. Pat. Nos. 4,399,216, 4,634,665 and 5,179,017, all by Axel et al.). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr- host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

10 *Non-Hybridoma Host Cells and Methods
of Recombinantly Producing Protein*

Nucleic acid molecules encoding the heavy chain or an antigen-binding portion thereof and/or the light chain or an antigen-binding portion thereof of an anti-IGF-IR antibody, and vectors comprising these nucleic acid molecules, can be used for transformation of a suitable mammalian host cell. Transformation can be by any known method for introducing polynucleotides into a host cell. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, biolistic injection and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be introduced into mammalian cells by viral vectors. Methods of transforming cells are well known in the art. See, e.g., U.S. Patent Nos. 4,399,216, 4,912,040, 4,740,461, and 4,959,455 (which patents are hereby incorporated herein by reference).

Mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC). These include, *inter alia*, Chinese hamster ovary (CHO) cells, NSO, SP2 cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), A549 cells, 3T3 cells, and a number of other cell lines. Mammalian host cells include human, mouse, rat, dog, monkey, pig, goat, bovine, horse and hamster cells. Cell lines of particular preference are selected through determining which cell lines have high expression levels. Other cell lines that may be used are insect cell lines, such as Sf9 cells, amphibian cells, bacterial cells, plant cells and fungal cells. When recombinant

expression vectors encoding the heavy chain or antigen-binding portion thereof, the light chain and/or antigen-binding portion thereof are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods.

Further, expression of antibodies of the invention (or other moieties therefrom) from production cell lines can be enhanced using a number of known techniques. For example, the glutamine synthetase gene expression system (the GS system) is a common approach for enhancing expression under certain conditions. The GS system is discussed in whole or part in connection with European Patent Nos. 0 216 846, 0 256 055, and 0 323 997 and European Patent Application No. 89303964.4.

It is likely that antibodies expressed by different cell lines or in transgenic animals will have different glycosylation from each other. However, all antibodies encoded by the nucleic acid molecules provided herein, or comprising the amino acid sequences provided herein are part of the instant invention, regardless of the glycosylation of the antibodies.

20 *Transgenic Animals*

The invention also provides transgenic non-human animals comprising one or more nucleic acid molecules of the invention that may be used to produce antibodies of the invention. Antibodies can be produced in and recovered from tissue or bodily fluids, such as milk, blood or urine, of goats, cows, horses, pigs, rats, mice, rabbits, hamsters or other mammals. See, e.g., U.S. Patent Nos. 5,827,690, 5,756,687, 5,750,172, and 5,741,957. As described above, non-human transgenic animals that comprise human immunoglobulin loci can be produced by immunizing with IGF-IR or a portion thereof.

In another embodiment, non-human transgenic animals are produced by introducing one or more nucleic acid molecules of the invention into the animal by standard transgenic techniques. See Hogan, *supra*. The transgenic cells used for making the transgenic animal can be embryonic stem cells or somatic cells. The transgenic non-human organisms can be chimeric, nonchimeric heterozygotes, and

nonchimeric homozygotes. See, e.g., Hogan *et al.*, Manipulating the Mouse Embryo: A Laboratory Manual 2ed., Cold Spring Harbor Press (1999); Jackson *et al.*, Mose Genetics and Transgenics: A Practical Approach, Oxford University Press (2000); and Pinkert, Transgenic Animal Technology: A Laboratory Handbook, Academic Press (1999). In another embodiment, the transgenic non-human organisms may have a targeted disruption and replacement that encodes a heavy chain and/or a light chain of interest. In a preferred embodiment, the transgenic animals comprise and express nucleic acid molecules encoding heavy and light chains that bind specifically to IGF-IR, preferably human IGF-IR. In another embodiment, the transgenic animals comprise nucleic acid molecules encoding a modified antibody such as a single-chain antibody, a chimeric antibody or a humanized antibody. The anti-IGF-IR antibodies may be made in any transgenic animal. In a preferred embodiment, the non-human animals are mice, rats, sheep, pigs, goats, cattle or horses. The non-human transgenic animal expresses said encoded polypeptides in blood, milk, urine, saliva, tears, mucus and other bodily fluids.

Phage Display Libraries

The invention provides a method for producing an anti-IGF-IR antibody or antigen-binding portion thereof comprising the steps of synthesizing a library of human antibodies on phage, screening the library with a IGF-IR or a portion thereof, isolating phage that bind IGF-IR, and obtaining the antibody from the phage. One method to prepare the library of antibodies comprises the steps of immunizing a non-human host animal comprising a human immunoglobulin locus with IGF-IR or an antigenic portion thereof to create an immune response, extracting cells from the host animal the cells that are responsible for production of antibodies; isolating RNA from the extracted cells, reverse transcribing the RNA to produce cDNA, amplifying the cDNA using a primer, and inserting the cDNA into phage display vector such that antibodies are expressed on the phage. Recombinant anti-IGF-IR antibodies of the invention may be obtained in this way.

Recombinant anti-IGF-IR human antibodies of the invention in addition to the anti-IGF-IR antibodies disclosed herein can be isolated by screening of a recombinant combinatorial antibody library, preferably a scFv phage display library, prepared using human VL and VH cDNAs prepared from mRNA derived from human lymphocytes. Methodologies for preparing and screening such libraries are known in

the art. There are commercially available kits for generating phage display libraries (e.g., the Pharmacia Recombinant Phage Antibody System, catalog no. 27-9400-01; and the Stratagene SurfZAP™ phage display kit, catalog no. 240612). There are also other methods and reagents that can be used in generating and screening antibody display libraries (see, e.g., Ladner et al. U.S. Pat. No. 5,223,409; Kang et al. PCT Publication No. WO 92/18619; Dower et al. PCT Publication No. WO 91/17271; Winter et al. PCT Publication No. WO 92/20791; Markland et al. PCT Publication No. WO 92/15679; Breitling et al. PCT Publication No. WO 93/01288; McCafferty et al. PCT Publication No. WO 92/01047; Garrard et al. PCT Publication No. WO 92/09690; Fuchs et al. (1991) Bio/Technology 9:1370-1372; Hay et al. (1992) Hum. Antibod. Hybridomas 3:81-85; Huse et al. (1989) Science 246:1275-1281; McCafferty et al., Nature (1990) 348:552-554; Griffiths et al. (1993) EMBO J 12:725-734; Hawkins et al. (1992) J. Mol. Biol. 226:889-896; Clackson et al. (1991) Nature 352:624-628; Gram et al. (1992) Proc. Natl. Acad. Sci. USA 89:3576-3580; Garrad et al. (1991) Bio/Technology 9:1373-1377; Hoogenboom et al. (1991) Nuc Acid Res 19:4133-4137; and Barbas et al. (1991) Proc. Natl. Acad. Sci. USA 88:7978-7982.

In a preferred embodiment, to isolate human anti-IGF-IR antibodies with the desired characteristics, a human anti-IGF-IR antibody as described herein is first used to select human heavy and light chain sequences having similar binding activity toward IGF-IR, using the epitope imprinting methods described in Hoogenboom et al., PCT Publication No. WO 93/06213. The antibody libraries used in this method are preferably scFv libraries prepared and screened as described in McCafferty et al., PCT Publication No. WO 92/01047, McCafferty et al., Nature (1990) 348:552-554; and Griffiths et al., (1993) EMBO J 12:725-734. The scFv antibody libraries preferably are screened using human IGF-IR as the antigen.

Once initial human VL and VH segments are selected, "mix and match" experiments, in which different pairs of the initially selected VL and VH segments are screened for IGF-IR binding, are performed to select preferred VL/VH pair combinations. Additionally, to further improve the quality of the antibody, the VL and VH segments of the preferred VL/VH pair(s) can be randomly mutated, preferably within the CDR3 region of VH and/or VL, in a process analogous to the *in vivo* somatic mutation process responsible for affinity maturation of antibodies during

a natural immune response. This *in vitro* affinity maturation can be accomplished by amplifying VH and VL regions using PCR primers complimentary to the VH CDR3 or VL CDR3, respectively, which primers have been "spiked" with a random mixture of the four nucleotide bases at certain positions such that the resultant PCR products
5 encode VH and VL segments into which random mutations have been introduced into the VH and/or VL CDR3 regions. These randomly mutated VH and VL segments can be rescreened for binding to IGF-IR.

Following screening and isolation of an anti-IGF-IR antibody of the invention from a recombinant immunoglobulin display library, nucleic acid encoding
10 the selected antibody can be recovered from the display package (e.g., from the phage genome) and subcloned into other expression vectors by standard recombinant DNA techniques. If desired, the nucleic acid can be further manipulated to create other antibody forms of the invention, as described below. To express a recombinant human antibody isolated by screening of a combinatorial library, the DNA encoding
15 the antibody is cloned into a recombinant expression vector and introduced into a mammalian host cells, as described above.

Class Switching

Another aspect of the instant invention is to provide a mechanism by which the class of an anti-IGF-IR antibody may be switched with another. In one
20 aspect of the invention, a nucleic acid molecule encoding VL or VH is isolated using methods well-known in the art such that it does not include any nucleic acid sequences encoding CL or CH. The nucleic acid molecule encoding VL or VH are then operatively linked to a nucleic acid sequence encoding a CL or CH from a different class of immunoglobulin molecule. This may be achieved using a vector or
25 nucleic acid molecule that comprises a CL or CH chain, as described above. For example, an anti-IGF-IR antibody that was originally IgM may be class switched to an IgG. Further, the class switching may be used to convert one IgG subclass to another, e.g., from IgG1 to IgG2. A preferred method for producing an antibody of the invention comprising a desired isotypes comprises the steps of isolating a nucleic
30 acid encoding the heavy chain of an anti-IGF-IR antibody and a nucleic acid encoding the light chain of an anti-IGF-IR antibody, obtaining the variable region of the heavy chain, ligating the variable region of the heavy chain with the constant domain of a

heavy chain of the desired isotype, expressing the light chain and the ligated heavy chain in a cell, and collecting the anti-IGF-IR antibody with the desired isotype.

Antibody Derivatives

One may use the nucleic acid molecules described above to generate antibody derivatives using techniques and methods known to one of ordinary skill in the art.

Humanized Antibodies

As was discussed above in connection with human antibody generation, there are advantages to producing antibodies with reduced immunogenicity. This can be accomplished to some extent using techniques of humanization and display techniques using appropriate libraries. It will be appreciated that murine antibodies or antibodies from other species can be humanized or primatized using techniques well known in the art. *See e.g., Winter and Harris Immunol Today* 14:43-46 (1993) and Wright et al. *Crit. Reviews in Immunol.* 12:125-168 (1992). The antibody of interest may be engineered by recombinant DNA techniques to substitute the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence (*see* WO 92/02190 and U.S. Patent Nos. 5,530,101, 5,585,089, 5,693,761, 5,693,792, 5,714,350, and 5,777,085). In a preferred embodiment, the anti-IGF-IR antibody can be humanized by substituting the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence while maintaining all of the CDRS of the heavy chain, the light chain or both the heavy and light chains.

Mutated Antibodies

In another embodiment, the nucleic acid molecules, vectors and host cells may be used to make mutated anti-IGF-IR antibodies. The antibodies may be mutated in the variable domains of the heavy and/or light chains to alter a binding property of the antibody. For example, a mutation may be made in one or more of the CDR regions to increase or decrease the K_d of the antibody for IGF-IR, to increase or decrease K_{off} , or to alter the binding specificity of the antibody. Techniques in site-directed mutagenesis are well-known in the art. *See, e.g., Sambrook et al. and Ausubel et al., supra.* In a preferred embodiment, mutations are made at an amino

acid residue that is known to be changed compared to germline in a variable region of an anti-IGF-IR antibody. In a more preferred embodiment, one or more mutations are made at an amino acid residue that is known to be changed compared to the germline in a variable region or CDR region of one of the anti-IGF-IR antibodies 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another embodiment, one or more mutations are made at an amino acid residue that is known to be changed compared to the germline in a variable region or CDR region whose amino acid sequence is presented in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24, or whose nucleic acid sequence is presented in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21 or 23. In another embodiment, the nucleic acid molecules are mutated in one or more of the framework regions. A mutation may be made in a framework region or constant domain to increase the half-life of the anti-IGF-IR antibody. See, e.g., WO 00/09560, published February 24, 2000, herein incorporated by reference. In one embodiment, there may be one, three or five point mutations and no more than ten point mutations. A mutation in a framework region or constant domain may also be made to alter the immunogenicity of the antibody, to provide a site for covalent or non-covalent binding to another molecule, or to alter such properties as complement fixation. Mutations may be made in each of the framework regions, the constant domain and the variable regions in a single mutated antibody. Alternatively, mutations may be made in only one of the framework regions, the variable regions or the constant domain in a single mutated antibody.

In one embodiment, there are no greater than ten amino acid changes in either the VH or VL regions of the mutated anti-IGF-IR antibody compared to the anti-IGF-IR antibody prior to mutation. In a more preferred embodiment, there is no more than five amino acid changes in either the VH or VL regions of the mutated anti-IGF-IR antibody, more preferably no more than three amino acid changes. In another embodiment, there are no more than fifteen amino acid changes in the constant domains, more preferably, no more than ten amino acid changes, even more preferably, no more than five amino acid changes.

30 *Modified Antibodies*

In another embodiment, a fusion antibody or immunoadhesin may be made which comprises all or a portion of an anti-IGF-IR antibody linked to another polypeptide. In a preferred embodiment, only the variable regions of the anti-IGF-IR

antibody are linked to the polypeptide. In another preferred embodiment, the VH domain of an anti-IGF-IR antibody are linked to a first polypeptide, while the VL domain of an anti-IGF-IR antibody are linked to a second polypeptide that associates with the first polypeptide in a manner in which the VH and VL domains can interact with one another to form an antibody binding site. In another preferred embodiment, the VH domain is separated from the VL domain by a linker such that the VH and VL domains can interact with one another (see below under Single Chain Antibodies). The VH-linker-VL antibody is then linked to the polypeptide of interest. The fusion antibody is useful to directing a polypeptide to an IGF-IR-expressing cell or tissue.

10 The polypeptide may be a therapeutic agent, such as a toxin, growth factor or other regulatory protein, or may be a diagnostic agent, such as an enzyme that may be easily visualized, such as horseradish peroxidase. In addition, fusion antibodies can be created in which two (or more) single-chain antibodies are linked to one another. This is useful if one wants to create a divalent or polyvalent antibody on a single

15 polypeptide chain, or if one wants to create a bispecific antibody.

To create a single chain antibody, (scFv) the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly₄-Ser)₃ (SEQ ID NO: 60), such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the

20 VL and VH regions joined by the flexible linker (see e.g., Bird et al. (1988) *Science* 242:423-426; Huston et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883; McCafferty et al., *Nature* (1990) 348:552-554). The single chain antibody may be monovalent, if only a single VH and VL are used, bivalent, if two VH and VL are used, or polyvalent, if more than two VH and VL are used.

25 In another embodiment, other modified antibodies may be prepared using anti-IGF-IR-encoding nucleic acid molecules. For instance, "Kappa bodies" (Ill et al., *Protein Eng* 10: 949-57 (1997)), "Minibodies" (Martin et al., *EMBO J* 13: 5303-9 (1994)), "Diabodies" (Holliger et al., *PNAS USA* 90: 6444-6448 (1993)), or "Janusins" (Traunecker et al., *EMBO J* 10: 3655-3659 (1991) and Traunecker et al.

30 "Janusin: new molecular design for bispecific reagents" *Int J Cancer Suppl* 7:51-52 (1992)) may be prepared using standard molecular biological techniques following the teachings of the specification.

In another aspect, chimeric and bispecific antibodies can be generated. A chimeric antibody may be made that comprises CDRs and framework regions from different antibodies. In a preferred embodiment, the CDRs of the chimeric antibody comprises all of the CDRs of the variable region of a light chain or heavy chain of an anti-IGF-IR antibody, while the framework regions are derived from one or more different antibodies. In a more preferred embodiment, the CDRs of the chimeric antibody comprise all of the CDRs of the variable regions of the light chain and the heavy chain of an anti-IGF-IR antibody. The framework regions may be from another species and may, in a preferred embodiment, be humanized. Alternatively, the framework regions may be from another human antibody.

A bispecific antibody can be generated that binds specifically to IGF-IR through one binding domain and to a second molecule through a second binding domain. The bispecific antibody can be produced through recombinant molecular biological techniques, or may be physically conjugated together. In addition, a single chain antibody containing more than one VH and VL may be generated that binds specifically to IGF-IR and to another molecule. Such bispecific antibodies can be generated using techniques that are well known for example, in connection with (i) and (ii) *see e.g.*, Fanger et al. *Immunol Methods* 4: 72-81 (1994) and Wright and Harris, *supra.* and in connection with (iii) *see e.g.*, Traunecker et al. *Int. J. Cancer (Suppl.)* 7: 51-52 (1992). In a preferred embodiment, the bispecific antibody binds to IGF-IR and to another molecule expressed at high level on cancer or tumor cells. In a more preferred embodiment, the other molecule is erbB2 receptor, VEGF, CD20 or EGF-R.

In an embodiment, the modified antibodies described above are prepared using one or more of the variable regions or one or more CDR regions from one of the antibodies selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 or 6.1.1. In another embodiment, the modified antibodies are prepared using one or more of the variable regions or one or more CDR regions whose amino acid sequence is presented in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22 or 24, or whose nucleic acid sequence is presented in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21 or 23.

Derivatized and Labeled Antibodies

An antibody or antibody portion of the invention can be derivatized or linked to another molecule (e.g., another peptide or protein). In general, the

antibodies or portion thereof is derivatized such that the IGF-IR binding is not affected adversely by the derivatization or labeling. Accordingly, the antibodies and antibody portions of the invention are intended to include both intact and modified forms of the human anti-IGF-IR antibodies described herein. For example, an antibody or antibody portion of the invention can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (e.g., a bispecific antibody or a diabody), a detection agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

One type of derivatized antibody is produced by crosslinking two or more antibodies (of the same type or of different types, e.g., to create bispecific antibodies). Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (e.g., m-maleimidobenzoyl-N-hydroxysuccinimide ester) or homobifunctional (e.g., disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, Ill.

Another type of derivatized antibody is a labeled antibody. Useful detection agents with which an antibody or antibody portion of the invention may be derivatized include fluorescent compounds, including fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin, lanthanide phosphors and the like. An antibody may also be labeled with enzymes that are useful for detection, such as horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase, glucose oxidase and the like. When an antibody is labeled with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a reaction product that can be discerned. For example, when the agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody may also be labeled with biotin, and detected through indirect measurement of avidin or streptavidin binding. An antibody may be labeled with a magnetic agent, such as gadolinium. An antibody may also be labeled with a predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains,

epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

An anti-IGF-IR antibody may also be labeled with a radiolabeled amino acid. The radiolabel may be used for both diagnostic and therapeutic purposes.

5 For instance, the radiolabel may be used to detect IGF-IR-expressing tumors by x-ray or other diagnostic techniques. Further, the radiolabel may be used therapeutically as a toxin for cancerous cells or tumors. Examples of labels for polypeptides include, but are not limited to, the following radioisotopes or radionuclides -- ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I .

10 An anti-IGF-IR antibody may also be derivatized with a chemical group such as polyethylene glycol (PEG), a methyl or ethyl group, or a carbohydrate group. These groups may be useful to improve the biological characteristics of the antibody, e.g., to increase serum half-life or to increase tissue binding.

Pharmaceutical Compositions and Kits

15 The invention also relates to a pharmaceutical composition for the treatment of a hyperproliferative disorder in a mammal which comprises a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. In one embodiment, said pharmaceutical composition is for the treatment of cancer such as brain, lung, squamous cell, bladder, 20 gastric, pancreatic, breast, head, neck, renal, kidney, ovarian, prostate, colorectal, esophageal, gynecological or thyroid cancer. In another embodiment, said pharmaceutical composition relates to non-cancerous hyperproliferative disorders such as, without limitation, restenosis after angioplasty and psoriasis. In another embodiment, the invention relates to pharmaceutical compositions for the treatment of 25 a mammal that requires activation of IGF-IR, wherein the pharmaceutical composition comprises a therapeutically effective amount of an activating antibody of the invention and a pharmaceutically acceptable carrier. Pharmaceutical compositions comprising activating antibodies may be used to treat animals that lack sufficient IGF-I or IGF-II, or may be used to treat osteoporosis, frailty or disorders in which the 30 mammal secretes too little active growth hormone or is unable to respond to growth hormone.

The anti-IGF-IR antibodies of the invention can be incorporated into pharmaceutical compositions suitable for administration to a subject. Typically, the

pharmaceutical composition comprises an antibody of the invention and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Examples of pharmaceutically acceptable carriers include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Pharmaceutically acceptable substances such as wetting or minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody or antibody portion.

The compositions of this invention may be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for passive immunization of humans with other antibodies. The preferred mode of administration is parenteral (e.g., intravenous, subcutaneous, intraperitoneal, intramuscular). In a preferred embodiment, the antibody is administered by intravenous infusion or injection. In another preferred embodiment, the antibody is administered by intramuscular or subcutaneous injection.

Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the anti-IGF-IR antibody in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of

preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size
5 in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

The antibodies of the present invention can be administered by a variety of methods known in the art, although for many therapeutic applications, the
10 preferred route/mode of administration is intraperitoneal, subcutaneous, intramuscular, intravenous or infusion. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In one embodiment, the antibodies of the present invention can be administered as a single dose or may be administered as multiple doses.

15 In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters,
20 and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, e.g., Sustained and Controlled Release Drug Delivery Systems, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

In certain embodiments, the anti-IGF-IR of the invention may be orally
25 administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches,
30 capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the invention by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

Supplementary active compounds can also be incorporated into the compositions. In certain embodiments, an anti-IGF-IR of the invention is coformulated with and/or coadministered with one or more additional therapeutic agents, such as a chemotherapeutic agent, an antineoplastic agent or an anti-tumor agent. For example, an anti-IGF-IR antibody may be coformulated and/or coadministered with one or more additional therapeutic agents. These agents include, without limitation, antibodies that bind other targets (e.g., antibodies that bind one or more growth factors or cytokines, their cell surface receptors or IGF-I), IGF-I binding proteins, antineoplastic agents, chemotherapeutic agents, anti-tumor agents, antisense oligonucleotides against IGF-IR or IGF-I, peptide analogues that block IGF-IR activation, soluble IGF-IR, and/or one or more chemical agents that inhibit IGF-I production or activity, which are known in the art, e.g., octreotide. For a pharmaceutical composition comprising an activating antibody, the anti-IGF-IR antibody may be formulated with a factor that increases cell proliferation or prevents apoptosis. Such factors include growth factors such as IGF-I, and/or analogues of IGF-I that activate IGF-IR. Such combination therapies may require lower dosages of the anti-IGF-IR antibody as well as the co-administered agents, thus avoiding possible toxicities or complications associated with the various monotherapies. In one embodiment, the antibody and one or more additional therapeutic agent.

The pharmaceutical compositions of the invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of an antibody or antibody portion of the invention. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the antibody or antibody portion may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

Dosage regimens may be adjusted to provide the optimum desired response (e.g., a therapeutic or prophylactic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Pharmaceutical composition comprising the antibody or comprising a combination therapy comprising the antibody and one or more additional therapeutic agents may be formulated for single or multiple doses. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals. A particularly useful formulation is 5 mg/ml anti-IGF-IR antibody in a buffer of 20mM sodium citrate, pH 5.5, 140mM NaCl, and 0.2mg/ml polysorbate 80.

An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of an antibody or antibody portion of the invention is 0.1-100 mg/kg, more preferably 0.5-50 mg/kg, more preferably 1-20 mg/kg, and even more preferably 1-10 mg/kg. It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. In one embodiment, the therapeutically or prophylactically effective amount of an antibody or antigen-binding portion thereof is administered along with one or more additional therapeutic agents.

In another aspect, the invention relates to administration of an anti-IGF-IR antibody for the treatment of cancer in a dose of less than 300 mg per month.

Another aspect of the present invention provides kits comprising the anti-IGF-IR antibodies and the pharmaceutical compositions comprising these antibodies. A kit may include, in addition to the antibody or pharmaceutical composition, diagnostic or therapeutic agents. A kit may also include instructions for
5 use in a diagnostic or therapeutic method. In a preferred embodiment, the kit includes the antibody or a pharmaceutical composition thereof and a diagnostic agent that can be used in a method described below. In another preferred embodiment, the kit includes the antibody or a pharmaceutical composition thereof and one or more therapeutic agents, such as an additional antineoplastic agent, anti-tumor agent or
10 chemotherapeutic agent, that can be used in a method described below.

This invention also relates to pharmaceutical compositions for inhibiting abnormal cell growth in a mammal which comprise an amount of a compound of the invention in combination with an amount of a chemotherapeutic agent, wherein the amounts of the compound, salt, solvate, or prodrug, and of the
15 chemotherapeutic agent are together effective in inhibiting abnormal cell growth. Many chemotherapeutic agents are presently known in the art. In one embodiment, the chemotherapeutic agents is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, anti-survival
20 agents, biological response modifiers, anti-hormones, e.g. anti-androgens, and anti-angiogenesis agents.

Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-II (cyclooxygenase II) inhibitors, can be used in conjunction with a compound of the
25 invention. Examples of useful COX-II inhibitors include CELEBREX™ (alecoxib), valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published October 24, 1996), WO 96/27583 (published March 7, 1996), European Patent Application No. 97304971.1 (filed July 8, 1997), European Patent Application No. 99308617.2 (filed October 29, 1999), WO
30 98/07697 (published February 26, 1998), WO 98/03516 (published January 29, 1998), WO 98/34918 (published August 13, 1998), WO 98/34915 (published August 13, 1998), WO 98/33768 (published August 6, 1998), WO 98/30566 (published July 16, 1998), European Patent Publication 606,046 (published July 13, 1994), European

Patent Publication 931,788 (published July 28, 1999), WO 90/05719 (published May 31, 1990), WO 99/52910 (published October 21, 1999), WO 99/52889 (published October 21, 1999), WO 99/29667 (published June 17, 1999), PCT International Application No. PCT/TB98/01113 (filed July 21, 1998), European Patent Application No. 99302232.1 (filed March 25, 1999), Great Britain patent application number 9912961.1 (filed June 3, 1999), United States Provisional Application No. 60/148,464 (filed August 12, 1999), United States Patent 5,863,949 (issued January 26, 1999), United States Patent 5,861,510 (issued January 19, 1999), and European Patent Publication 780,386 (published June 25, 1997), all of which are incorporated herein in their entireties by reference. Preferred MMP inhibitors are those that do not demonstrate arthralgia. More preferred, are those that selectively inhibit MMP-2 and/or MMP-9 relative to the other matrix-metalloproteinases (i.e. MMP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13). Some specific examples of MMP inhibitors useful in the present invention are AG-3340, RO 32-3555, RS 13-0830, and the compounds recited in the following list: 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclopentyl)-amino]-propionic acid; 3-exo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; (2R, 3R) 1-[4-(2-chloro-4-fluoro-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide; 4-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclobutyl)-amino]-propionic acid; 4-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide; (R) 3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-3-carboxylic acid hydroxyamide; (2R, 3R) 1-[4-(4-fluoro-2-methyl-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-1-methyl-ethyl)-amino]-propionic acid; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(4-hydroxycarbamoyl-tetrahydro-pyran-4-yl)-amino]-propionic acid; 3-exo-3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-8-oxa-icyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; 3-endo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-icyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; and (R) 3-[4-(4-fluoro-phenoxy)-

benzenesulfonylamino]-tetrahydro-furan-3-carboxylic acid hydroxyamide; and pharmaceutically acceptable salts and solvates of said compounds.

A compound of the invention can also be used with signal transduction inhibitors, such as agents that can inhibit EGF-R (epidermal growth factor receptor) responses, such as EGF-R antibodies, EGF antibodies, and molecules that are EGF-R inhibitors; VEGF (vascular endothelial growth factor) inhibitors, such as VEGF receptors and molecules that can inhibit VEGF; and erbB2 receptor inhibitors, such as organic molecules or antibodies that bind to the erbB2 receptor, for example, HERCEPTINTM (Genentech, Inc.). EGF-R inhibitors are described in, for example in WO 95/19970 (published July 27, 1995), WO 98/14451 (published April 9, 1998), WO 98/02434 (published January 22, 1998), and United States Patent 5,747,498 (issued May 5, 1998), and such substances can be used in the present invention as described herein. EGFR-inhibiting agents include, but are not limited to, the monoclonal antibodies C225 and anti-EGFR 22Mab (ImClone Systems Incorporated), ABX-EGF (Abgenix/Cell Genesys), EMD-7200 (Merck KgaA), EMD-5590 (Merck KgaA), MDX-447/H-477 (Medarex Inc. and Merck KgaA), and the compounds ZD-1834, ZD-1838 and ZD-1839 (AstraZeneca), PKI-166 (Novartis), PKI-166/CGP-75166 (Novartis), PTK 787 (Novartis), CP 701 (Cephalon), leflunomide (Pharmacia/Sugen), CI-1033 (Warner Lambert Parke Davis), CI-1033/PD 183,805 (Warner Lambert Parke Davis), CL-387,785 (Wyeth-Ayerst), BBR-1611 (Boehringer Mannheim GmbH/Roche), Naamidine A (Bristol Myers Squibb), RC-3940-II (Pharmacia), BIBX-1382 (Boehringer Ingelheim), OLX-103 (Merck & Co.), VRCTC-310 (Ventech Research), EGF fusion toxin (Seragen Inc.), DAB-389 (Seragen/Lilgand), ZM-252808 (Imperial Cancer Research Fund), RG-50864 (INSERM), LFM-A12 (Parker Hughes Cancer Center), WHI-P97 (Parker Hughes Cancer Center), GW-282974 (Glaxo), KT-8391 (Kyowa Hakko) and EGF-R Vaccine (York Medical/Centro de Immunologia Molecular (CIM)). These and other EGF-R-inhibiting agents can be used in the present invention.

VEGF inhibitors, for example SU-5416 and SU-6668 (Sugen Inc.), SH-268 (Schering), and NX-1838 (NeXstar) can also be combined with the compound of the present invention. VEGF inhibitors are described in, for example in WO 99/24440 (published May 20, 1999), PCT International Application PCT/IB99/00797 (filed May 3, 1999), in WO 95/21613 (published August 17, 1995),

WO 99/61422 (published December 2, 1999), United States Patent 5,834,504 (issued November 10, 1998), WO 98/50356 (published November 12, 1998), United States Patent 5,883,113 (issued March 16, 1999), United States Patent 5,886,020 (issued March 23, 1999), United States Patent 5,792,783 (issued August 11, 1998), WO 5 99/10349 (published March 4, 1999), WO 97/32856 (published September 12, 1997), WO 97/22596 (published June 26, 1997), WO 98/54093 (published December 3, 1998), WO 98/02438 (published January 22, 1998), WO 99/16755 (published April 8, 1999), and WO 98/02437 (published January 22, 1998), all of which are incorporated herein in their entireties by reference. Other examples of some specific VEGF 10 inhibitors useful in the present invention are IM862 (Cytran Inc.); anti-VEGF monoclonal antibody of Genentech, Inc.; and angiozyme, a synthetic ribozyme from Ribozyme and Chiron. These and other VEGF inhibitors can be used in the present invention as described herein.

ErbB2 receptor inhibitors, such as GW-282974 (Glaxo Wellcome plc), 15 and the monoclonal antibodies AR-209 (Aronex Pharmaceuticals Inc.) and 2B-1 (Chiron), can furthermore be combined with the compound of the invention, for example those indicated in WO 98/02434 (published January 22, 1998), WO 99/35146 (published July 15, 1999), WO 99/35132 (published July 15, 1999), WO 98/02437 (published January 22, 1998), WO 97/13760 (published April 17, 1997), 20 WO 95/19970 (published July 27, 1995), United States Patent 5,587,458 (issued December 24, 1996), and United States Patent 5,877,305 (issued March 2, 1999), which are all hereby incorporated herein in their entireties by reference. ErbB2 receptor inhibitors useful in the present invention are also described in United States Provisional Application No. 60/117,341, filed January 27, 1999, and in United States 25 Provisional Application No. 60/117,346, filed January 27, 1999, both of which are incorporated in their entireties herein by reference. The erbB2 receptor inhibitor compounds and substance described in the aforementioned PCT applications, U.S. patents, and U.S. provisional applications, as well as other compounds and substances that inhibit the erbB2 receptor, can be used with the compound of the present 30 invention in accordance with the present invention.

Anti-survival agents include anti-IGF-IR antibodies and anti-integrin agents, such as anti-integrin antibodies.

Diagnostic Methods of Use

The anti-IGF-IR antibodies may be used to detect IGF-IR in a biological sample *in vitro* or *in vivo*. The anti-IGF-IR antibodies may be used in a conventional immunoassay, including, without limitation, an ELISA, an RIA, FACS, tissue immunohistochemistry, Western blot or immunoprecipitation. The anti-IGF-IR antibodies of the invention may be used to detect IGF-IR from humans. In another embodiment, the anti-IGF-IR antibodies may be used to detect IGF-IR from Old World primates such as cynomologous and rhesus monkeys, chimpanzees and apes. The invention provides a method for detecting anti-IGF-IR in a biological sample comprising contacting a biological sample with an anti-IGF-IR antibody of the invention and detecting the bound antibody bound to anti-IGF-IR, to detect the IGF-IR in the biological sample. In one embodiment, the anti-IGF-IR antibody is directly labeled with a detectable label. In another embodiment, the anti-IGF-IR antibody (the first antibody) is unlabeled and a second antibody or other molecule that can bind the anti-IGF-IR antibody is labeled. As is well known to one of skill in the art, a second antibody is chosen that is able to specifically bind the specific species and class of the first antibody. For example, if the anti-IGF-IR antibody is a human IgG, then the secondary antibody may be an anti-human-IgG. Other molecules that can bind to antibodies include, without limitation, Protein A and Protein G, both of which are available commercially, e.g., from Pierce Chemical Co.

Suitable labels for the antibody or secondary have been disclosed *supra*, and include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, magnetic agents and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; an example of a magnetic agent includes gadolinium; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

In an alternative embodiment, IGF-IR can be assayed in a biological sample by a competition immunoassay utilizing IGF-IR standards labeled with a

detectable substance and an unlabeled anti-IGF-IR antibody. In this assay, the biological sample, the labeled IGF-IR standards and the anti-IGF-IR antibody are combined and the amount of labeled IGF-IR standard bound to the unlabeled antibody is determined. The amount of IGF-IR in the biological sample is inversely
5 proportional to the amount of labeled IGF-IR standard bound to the anti-IGF-IR antibody.

One may use the immunoassays disclosed above for a number of purposes. In one embodiment, the anti-IGF-IR antibodies may be used to detect IGF-IR in cells in cell culture. In a preferred embodiment, the anti-IGF-IR antibodies may
10 be used to determine the level of tyrosine phosphorylation, tyrosine autophosphorylation of IGF-IR, and/or the amount of IGF-IR on the cell surface after treatment of the cells with various compounds. This method can be used to test compounds that may be used to activate or inhibit IGF-IR. In this method, one sample of cells is treated with a test compound for a period of time while another
15 sample is left untreated. If tyrosine autophosphorylation is to be measured, the cells are lysed and tyrosine phosphorylation of the IGF-IR is measured using an immunoassay described above or as described in Example III, which uses an ELISA. If the total level of IGF-IR is to be measured, the cells are lysed and the total IGF-IR level is measured using one of the immunoassays described above.

20 A preferred immunoassay for determining IGF-IR tyrosine phosphorylation or for measuring total IGF-IR levels is an ELISA or Western blot. If only the cell surface level of IGF-IR is to be measured, the cells are not lysed, and the cell surface levels of IGF-IR are measured using one of the immunoassays described above. A preferred immunoassay for determining cell surface levels of IGF-IR
25 includes the steps of labeling the cell surface proteins with a detectable label, such as biotin or ^{125}I , immunoprecipitating the IGF-IR with an anti-IGF-IR antibody and then detecting the labeled IGF-IR. Another preferred immunoassay for determining the localization of IGF-IR, e.g., cell surface levels, is by using immunohistochemistry. Methods such as ELISA, RIA, Western blot, immunohistochemistry, cell surface
30 labeling of integral membrane proteins and immunoprecipitation are well known in the art. See, e.g., Harlow and Lane, *supra*. In addition, the immunoassays may be scaled up for high throughput screening in order to test a large number of compounds for either activation or inhibition of IGF-IR.

The anti-IGF-IR antibodies of the invention may also be used to determine the levels of IGF-IR in a tissue or in cells derived from the tissue. In a preferred embodiment, the tissue is a diseased tissue. In a more preferred embodiment, the tissue is a tumor or a biopsy thereof. In a preferred embodiment of the method, a tissue or a biopsy thereof is excised from a patient. The tissue or biopsy is then used in an immunoassay to determine, e.g., IGF-IR levels, cell surface levels of IGF-IR, levels of tyrosine phosphorylation of IGF-IR, or localization of IGF-IR by the methods discussed above. The method can be used to determine if a tumor expresses IGF-IR at a high level.

10 The above-described diagnostic method can be used to determine whether a tumor expresses high levels of IGF-IR, which may be indicative that the tumor will respond well to treatment with anti-IGF-IR antibody. The diagnostic method may also be used to determine whether a tumor is potentially cancerous, if it expresses high levels of IGF-IR, or benign, if it expresses low levels of IGF-IR.

15 Further, the diagnostic method may also be used to determine whether treatment with anti-IGF-IR antibody (see below) is causing a tumor to express lower levels of IGF-IR and/or to express lower levels of tyrosine autophosphorylation, and thus can be used to determine whether the treatment is successful. In general, a method to determine whether an anti-IGF-IR antibody decreases tyrosine phosphorylation

20 comprises the steps of measuring the level of tyrosine phosphorylation in a cell or tissue of interest, incubating the cell or tissue with an anti-IGF-IR antibody or antigen-binding portion thereof, then re-measuring the level of tyrosine phosphorylation in the cell or tissue. The tyrosine phosphorylation of IGF-IR or of another protein(s) may be measured. The diagnostic method may also be used to

25 determine whether a tissue or cell is not expressing high enough levels of IGF-IR or high enough levels of activated IGF-IR, which may be the case for individuals with dwarfism, osteoporosis or diabetes. A diagnosis that levels of IGF-IR or active IGF-IR are too low could be used for treatment with activating anti-IGF-IR antibodies, IGF-I or other therapeutic agents for increasing IGF-IR levels or activity.

30 The antibodies of the present invention may also be used *in vivo* to localize tissues and organs that express IGF-IR. In a preferred embodiment, the anti-IGF-IR antibodies can be used to localize IGF-IR-expressing tumors. The advantage of the anti-IGF-IR antibodies of the present invention is that they will not generate an

immune response upon administration. The method comprises the steps of administering an anti-IGF-IR antibody or a pharmaceutical composition thereof to a patient in need of such a diagnostic test and subjecting the patient to imaging analysis to determine the location of the IGF-IR-expressing tissues. Imaging analysis is well known in the medical art, and includes, without limitation, x-ray analysis, magnetic resonance imaging (MRI) or computed tomography (CE). In another embodiment of the method, a biopsy is obtained from the patient to determine whether the tissue of interest expresses IGF-IR rather than subjecting the patient to imaging analysis. In a preferred embodiment, the anti-IGF-IR antibodies may be labeled with a detectable agent that can be imaged in a patient. For example, the antibody may be labeled with a contrast agent, such as barium, which can be used for x-ray analysis, or a magnetic contrast agent, such as a gadolinium chelate, which can be used for MRI or CE. Other labeling agents include, without limitation, radioisotopes, such as ^{99}Tc . In another embodiment, the anti-IGF-IR antibody will be unlabeled and will be imaged by administering a second antibody or other molecule that is detectable and that can bind the anti-IGF-IR antibody.

Therapeutic Methods of Use

In another embodiment, the invention provides a method for inhibiting IGF-IR activity by administering an anti-IGF-IR antibody to a patient in need thereof. Any of the types of antibodies described herein may be used therapeutically. In a preferred embodiment, the anti-IGF-IR antibody is a human, chimeric or humanized antibody. In another preferred embodiment, the IGF-IR is human and the patient is a human patient. Alternatively, the patient may be a mammal that expresses an IGF-IR that the anti-IGF-IR antibody cross-reacts with. The antibody may be administered to a non-human mammal expressing an IGF-IR with which the antibody cross-reacts (i.e. a primate, or a cynomologous or rhesus monkey) for veterinary purposes or as an animal model of human disease. Such animal models may be useful for evaluating the therapeutic efficacy of antibodies of this invention.

As used herein, the term "a disorder in which IGF-IR activity is detrimental" is intended to include diseases and other disorders in which the presence of high levels of IGF-IR in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in

which high levels of IGF-IR activity is detrimental is a disorder in which inhibition of IGF-IR activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the levels of IGF-IR on the cell surface or in increased tyrosine autophosphorylation of IGF-IR in the affected cells or tissues of a subject suffering from the disorder. The increase in IGF-IR levels may be detected, for example, using an anti-IGF-IR antibody as described above.

In a preferred embodiment, an anti-IGF-IR antibody may be administered to a patient who has an IGF-IR-expressing tumor. A tumor may be a solid tumor or may be a non-solid tumor, such as a lymphoma. In a more preferred embodiment, an anti-IGF-IR antibody may be administered to a patient who has an IGF-IR-expressing tumor that is cancerous. In an even more preferred embodiment, the anti-IGF-IR antibody is administered to a patient who has a tumor of the lung, breast, prostate or colon. In a highly preferred embodiment, the method causes the tumor not to increase in weight or volume or to decrease in weight or volume. In another embodiment, the method causes the IGF-IR on the tumor to be internalized. In a preferred embodiment, the antibody is selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding region thereof.

In another preferred embodiment, an anti-IGF-IR antibody may be administered to a patient who expresses inappropriately high levels of IGF-I. It is known in the art that high-level expression of IGF-I can lead to a variety of common cancers. In a more preferred embodiment, the anti-IGF-IR antibody is administered to a patient with prostate cancer, glioma or fibrosarcoma. In an even more preferred embodiment, the method causes the cancer to stop proliferating abnormally, or not to increase in weight or volume or to decrease in weight or volume.

In one embodiment, said method relates to the treatment of cancer such as brain, squamous cell, bladder, gastric, pancreatic, breast, head, neck, esophageal, prostate, colorectal, lung, renal, kidney, ovarian, gynecological or thyroid cancer. Patients that can be treated with a compounds of the invention according to the methods of this invention include, for example, patients that have been diagnosed as having lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head and neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal

cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer, gynecologic tumors (e.g., uterine sarcomas, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina or carcinoma of the vulva), Hodgkin's disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system (e.g., cancer of the thyroid, parathyroid or adrenal glands), sarcomas of soft tissues, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemia, solid tumors of childhood, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter (e.g., renal cell carcinoma, carcinoma of the renal pelvis), or neoplasms of the central nervous system (e.g., primary CNS lymphoma, spinal axis tumors, brain stem gliomas or pituitary adenomas).

The antibody may be administered once, but more preferably is administered multiple times. The antibody may be administered from three times daily to once every six months. The administering may be on a schedule such as three times daily, twice daily, once daily, once every two days, once every three days, once weekly, once every two weeks, once every month, once every two months, once every three months and once every six months. The antibody may be administered via an oral, mucosal, buccal, intranasal, inhalable, intravenous, subcutaneous, intramuscular, parenteral, intratumor or topical route. The antibody may be administered at a site distant from the site of the tumor. The antibody may also be administered continuously via a minipump. The antibody may be administered once, at least twice or for at least the period of time until the condition is treated, palliated or cured. The antibody generally will be administered for as long as the tumor is present provided that the antibody causes the tumor or cancer to stop growing or to decrease in weight or volume. The antibody will generally be administered as part of a pharmaceutical composition as described *supra*. The dosage of antibody will generally be in the range of 0.1-100 mg/kg, more preferably 0.5-50 mg/kg, more preferably 1-20 mg/kg, and even more preferably 1-10 mg/kg. The serum concentration of the antibody may be measured by any method known in the art. See, e.g., Example XVII below. The antibody may also be administered prophylactically in order to prevent a cancer or tumor from occurring. This may be especially useful in patients that have a "high normal" level of IGF-I because these patients have been shown to have a higher risk of developing common cancers. See Rosen et al., *supra*.

In another aspect, the anti-IGF-IR antibody may be co-administered with other therapeutic agents, such as antineoplastic drugs or molecules, to a patient who has a hyperproliferative disorder, such as cancer or a tumor. In one aspect, the invention relates to a method for the treatment of the hyperproliferative disorder in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of the invention in combination with an anti-tumor agent selected from the group consisting of, but not limited to, mitotic inhibitors, alkylating agents, anti-metabolites, intercalating agents, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, kinase inhibitors, matrix metalloprotease inhibitors, genetic therapeutics and anti-androgens. In a more preferred embodiment, the antibody may be administered with an antineoplastic agent, such as adriamycin or taxol. In another preferred embodiment, the antibody or combination therapy is administered along with radiotherapy, chemotherapy, photodynamic therapy, surgery or other immunotherapy. In yet another preferred embodiment, the antibody will be administered with another antibody. For example, the anti-IGF-IR antibody may be administered with an antibody or other agent that is known to inhibit tumor or cancer cell proliferation, e.g., an antibody or agent that inhibits erbB2 receptor, EGF-R, CD20 or VEGF.

Co-administration of the antibody with an additional therapeutic agent (combination therapy) encompasses administering a pharmaceutical composition comprising the anti-IGF-IR antibody and the additional therapeutic agent and administering two or more separate pharmaceutical compositions, one comprising the anti-IGF-IR antibody and the other(s) comprising the additional therapeutic agent(s). Further, although co-administration or combination therapy generally means that the antibody and additional therapeutic agents are administered at the same time as one another, it also encompasses instances in which the antibody and additional therapeutic agents are administered at different times. For instance, the antibody may be administered once every three days, while the additional therapeutic agent is administered once daily. Alternatively, the antibody may be administered prior to or subsequent to treatment of the disorder with the additional therapeutic agent. Similarly, administration of the anti-IGF-IR antibody may be administered prior to or subsequent to other therapy, such as radiotherapy, chemotherapy, photodynamic therapy, surgery or other immunotherapy.

The antibody and one or more additional therapeutic agents (the combination therapy) may be administered once, twice or at least the period of time until the condition is treated, palliated or cured. Preferably, the combination therapy is administered multiple times. The combination therapy may be administered from
5 three times daily to once every six months. The administering may be on a schedule such as three times daily, twice daily, once daily, once every two days, once every three days, once weekly, once every two weeks, once every month, once every two months, once every three months and once every six months, or may be administered continuously via a minipump. The combination therapy may be administered via an
10 oral, mucosal, buccal, intranasal, inhalable, intravenous, subcutaneous, intramuscular, parenteral, intratumor or topical route. The combination therapy may be administered at a site distant from the site of the tumor. The combination therapy generally will be administered for as long as the tumor is present provided that the antibody causes the tumor or cancer to stop growing or to decrease in weight or volume.

15 In a still further embodiment, the anti-IGF-IR antibody is labeled with a radiolabel, an immunotoxin or a toxin, or is a fusion protein comprising a toxic peptide. The anti-IGF-IR antibody or anti-IGF-IR antibody fusion protein directs the radiolabel, immunotoxin, toxin or toxic peptide to the IGF-IR-expressing tumor or cancer cell. In a preferred embodiment, the radiolabel, immunotoxin, toxin or toxic
20 peptide is internalized after the anti-IGF-IR antibody binds to the IGF-IR on the surface of the tumor or cancer cell.

In another aspect, the anti-IGF-IR antibody may be used therapeutically to induce apoptosis of specific cells in a patient in need thereof. In many cases, the cells targeted for apoptosis are cancerous or tumor cells. Thus, in a
25 preferred embodiment, the invention provides a method of inducing apoptosis by administering a therapeutically effective amount of an anti-IGF-IR antibody to a patient in need thereof. In a preferred embodiment, the antibody is selected from 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, or 6.1.1, or comprises a heavy chain, light chain or antigen-binding region thereof.

30 In another aspect, the anti-IGF-IR antibody may be used to treat noncancerous states in which high levels of IGF-I and/or IGF-IR have been associated with the noncancerous state or disease. In one embodiment, the method comprises the step of administering an anti-IGF-IR antibody to a patient who has a noncancerous

pathological state caused or exacerbated by high levels of IGF-I and/or IGF-IR levels or activity. In a preferred embodiment, the noncancerous pathological state is acromegaly, gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels or inappropriate microvascular proliferation, such as that found as a
5 complication of diabetes, especially of the eye. In a more preferred embodiment, the anti-IGF-IR antibody slows the progress of the noncancerous pathological state. In a more preferred embodiment, the anti-IGF-IR antibody stops or reverses, at least in part, the noncancerous pathological state.

In another aspect, the invention provides a method of administering an
10 activating anti-IGF-IR antibody to a patient in need thereof. In one embodiment, the activating antibody or pharmaceutical composition is administered to a patient in need thereof in an amount effective to increase IGF-IR activity. In a more preferred embodiment, the activating antibody is able to restore normal IGF-IR activity. In another preferred embodiment, the activating antibody may be administered to a
15 patient who has small stature, neuropathy, a decrease in muscle mass or osteoporosis. In another preferred embodiment, the activating antibody may be administered with one or more other factors that increase cell proliferation, prevent apoptosis or increase IGF-IR activity. Such factors include growth factors such as IGF-I, and/or analogues of IGF-I that activate IGF-IR. In a preferred embodiment, the antibody is selected
20 from 4.17.3, or comprises a heavy chain, light chain or antigen-binding portion thereof.

Gene Therapy

The nucleic acid molecules of the instant invention may be administered to a patient in need thereof via gene therapy. The therapy may be either
25 *in vivo* or *ex vivo*. In a preferred embodiment, nucleic acid molecules encoding both a heavy chain and a light chain are administered to a patient. In a more preferred embodiment, the nucleic acid molecules are administered such that they are stably integrated into the chromosome of B cells because these cells are specialized for producing antibodies. In a preferred embodiment, precursor B cells are transfected or
30 infected *ex vivo* and re-transplanted into a patient in need thereof. In another embodiment, precursor B cells or other cells are infected *in vivo* using a virus known to infect the cell type of interest. Typical vectors used for gene therapy include liposomes, plasmids, or viral vectors, such as retroviruses, adenoviruses and adeno-

associated viruses. After infection either *in vivo* or *ex vivo*, levels of antibody expression may be monitored by taking a sample from the treated patient and using any immunoassay known in the art and discussed herein.

In a preferred embodiment, the gene therapy method comprises the steps of administering an effective amount of an isolated nucleic acid molecule encoding the heavy chain or the antigen-binding portion thereof of the human antibody or portion thereof and expressing the nucleic acid molecule. In another embodiment, the gene therapy method comprises the steps of administering an effective amount of an isolated nucleic acid molecule encoding the light chain or the antigen-binding portion thereof of the human antibody or portion thereof and expressing the nucleic acid molecule. In a more preferred method, the gene therapy method comprises the steps of administering an effective amount of an isolated nucleic acid molecule encoding the heavy chain or the antigen-binding portion thereof of the human antibody or portion thereof and an effective amount of an isolated nucleic acid molecule encoding the light chain or the antigen-binding portion thereof of the human antibody or portion thereof and expressing the nucleic acid molecules. The gene therapy method may also comprise the step of administering another anti-cancer agent, such as taxol, tamoxifen, 5-FU, adriamycin or CP-358,774.

In order that this invention may be better understood, the following examples are set forth. These examples are for purposes of illustration only and are not to be construed as limiting the scope of the invention in any manner.

EXAMPLE I: Generation of Hybridomas Producing Anti-IGF-IR Antibody

Antibodies of the invention were prepared, selected, and assayed as follows:

Immunization and hybridoma generation

Eight to ten week old XENOMICE™ were immunized intraperitoneally or in their hind footpads with either the extracellular domain of human IGF-IR (10 µg/dose/mouse), or with 3T3-IGF-IR or 300.19-IGF-IR cells, which are two transfected cell lines that express human IGF-IR on their plasma membranes (10 x 10⁶ cells/dose/mouse). This dose was repeated five to seven times over a three to eight week period. Four days before fusion, the mice received a final injection of the extracellular domain of human IGF-IR in PBS. Spleen and lymph

node lymphocytes from immunized mice were fused with the non-secretory myeloma P3-X63-Ag8.653 cell line and were subjected to HAT selection as previously described (Galfre and Milstein, *Methods Enzymol.* 73:3-46, 1981). A panel of hybridomas all secreting IGF-IR specific human IgG2 κ antibodies were recovered.

- 5 Seven hybridomas producing monoclonal antibodies specific for IGF-IR were selected for further study and were designated 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2, 4.17.3 and 6.1.1.

Hybridomas 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3 were deposited in the American Type Culture Collection (ATCC), 10801 University
10 Boulevard, Manassas, VA 20110-2209, on December 12, 2000 with the following deposit numbers:

	<u>Hybridoma</u>	<u>Deposit No.</u>
	2.12.1	PTA-2792
	2.13.2	PTA-2788
15	2.14.3	PTA-2790
	3.1.1	PTA-2791
	4.9.2	PTA-2789
	4.17.3	PTA-2793

20 EXAMPLE II: Determination of Affinity Constants (K_d) of Fully Human Anti-IGF-IR Monoclonal Antibodies by BIAcore

We performed affinity measures of purified antibodies by surface plasmon resonance using the BIAcore 3000 instrument, following the manufacturer's protocols.

Protocol I

- 25 To perform kinetic analyses, protein-A was immobilized on the sensorchip surfaces of the BIAcore. The sensorchip was then used to capture the anti-IGF-IR antibodies of the present invention. Different concentrations of the extracellular domain of IGF-IR were injected on the sensorchip and the binding and dissociation kinetics of the interactions between the anti-IGF-IR antibodies and the
30 extracellular domain of IGF-IR were analyzed. The data were evaluated with global fit Langmuir 1:1, using baseline drift models available on the BIAevaluation software provided by BIAcore.

Protocol 2

BIAcore measurements were performed essentially as described by Fagerstam et al. "Detection of antigen-antibody interactions by surface plasmon resonance. Applications to epitope mapping." J. Mol. Recog. 3: 208-214. (1990).

- 5 Table I lists affinity measurements for representative anti-IGF-IR antibodies of the present invention:

Table I

Monoclonal Antibody	K_d (M) <i>Protocol 1</i>	K_d (M) <i>Protocol 2</i>
2.12.1	7.37×10^{-9}	
2.13.2	3.5×10^{-9}	1.53×10^{-9}
2.14.3	6.41×10^{-10}	
3.1.1	1.15×10^{-9}	
4.9.2	6.84×10^{-10}	4.27×10^{-10}
4.17.3	1.3×10^{-8}	
6.1.1	5.65×10^{-10}	

- The kinetic analyses indicates that the antibodies prepared in accordance with the invention possess high affinities and strong binding constants for the extracellular domain of IGF-IR.

EXAMPLE III: Antibody-mediated Inhibition of IGF-I-induced Phosphorylation of IGF-IR

- We performed ELISA experiments in order to determine whether the antibodies of this invention were able to block IGF-I-mediated activation of IGF-IR. IGF-I-mediated activation of IGF-IR was detected by increased receptor-associated tyrosine phosphorylation.

ELISA Plate Preparation

- We prepared ELISA capture plates by adding 100 μ l blocking buffer (3% bovine serum albumin [BSA] in Tris-buffered saline [TBS]) to each well of a ReactiBind Protein G-coated 96-well plates (Pierce) and incubated the plates with

shaking for 30 minutes at room temperature. We diluted rabbit pan-specific SC-713 anti-IGF-IR antibody (Santa Cruz) in blocking buffer to a concentration of 5 µg/ml and added 100 µl diluted antibody to each well. We incubated the plates with shaking for 60-90 minutes at room temperature. We then washed the plates five times with
5 wash buffer (TBS + 0.1% Tween 20) and gently blotted the remaining buffer out onto paper towels. These plates were not allowed to dry out prior to the addition of lysate.

Preparation of Lysate from IGF-IR-expressing Cells

We placed IGF-IR-transfected NIH-3T3 cells (5×10^4 /ml) in 100 µl of growth media (DMEM high glucose media supplemented with L-glutamine (0.29
10 mg/ml), 10% heat-inactivated FBS, and 500 µg/ml each of geneticin, penicillin and streptomycin) in 96-well U-bottom plates. We incubated the plates at 37°C, 5% CO₂ overnight to allow the cells to attach. We decanted the media from the plates and replaced it with 100 µl fresh growth media per well. For testing, we diluted the potential anti-IGF-IR antibodies to five times the desired final concentration in
15 growth media and added 25 µl per well. All samples were performed in triplicate. We then incubated the plates at 37°C for one hour. We stimulated the cells with 25 µl/well of 600 ng/ml IGF-1 (prepared in growth media) and incubate the plates at room temperature for 10 minutes. We then decanted the media by inverting the plates and blotting gently onto paper towels and lysed the adherent cells by adding 50 µl of
20 lysis buffer (50 mM HEPES, pH 7.4, 10 mM EDTA, 150 mM NaCl, 1.5 mM MgCl₂, 1.6 mM NaVO₄, 1% Triton X-100, 1% glycerol supplemented immediately before use with one EDTA-free protease inhibitor tablet [Roche Molecular Sciences] per 50 ml) and shaking for 5 minutes at room temperature. We added 200 µl dilution buffer (50 mM HEPES, pH 7.4, 1.6 mM NaVO₄) to each well and mixed by pipetting up and
25 down. We transferred 100 µl of lysate from each well to each well of the ELISA capture plate prepared as described above and incubated with gentle shaking for two hours at room temperature.

ELISA with Anti-tyrosine-phosphate (pTYR) Antibodies

We removed the cell lysate by inverting the plates, washed the plates
30 five times with wash buffer and blotted on paper towels. We added 100 µl per well pTYR-specific antibody (HRP-PY54) diluted in blocking buffer to a concentration of

0.2 µg/ml and incubated the plates with shaking for 30 minutes at room temperature. We then washed these plates five times with wash buffer and blotted on paper towels.

We detected binding of the HRP-PY54 antibody by adding 100 µl per well of TMB peroxidase substrate solution (Kirkegaard & Perry) and incubating with shaking as the color developed (approximately 2-10 minutes). We stopped the color development reaction by adding 100 µl per well of TMB stop solution (Kirkegaard & Perry). We then shook the plates for 10 seconds at room temperature to mix the solution and quantitated by measurement at OD_{450nm}.

Table II and Figure 4 show the results of this experiment performed with several antibodies of the invention. The results of this experiment demonstrate the ability of the antibodies of this invention to block IGF-I-mediated activation of IGF-IR as shown by increased receptor-associated tyrosine phosphorylation. Furthermore, these results can be used to quantify the relative potency of the antibodies of this invention.

Table II

Monoclonal Antibody	IC ₅₀ (µg/ml)
2.12.1	0.172
2.13.2	0.0812
2.14.3	0.325
4.9.2	0.0324

EXAMPLE IV: Antibody-mediated Blocking of IGF-I/IGF-IR Binding

We performed ELISA experiments to quantitate the ability of the antibodies of the invention to inhibit IGF-I binding to IGF-IR in a cell-based assay. We plated IGF-IR-transfected NIH-3T3 cells (5x10⁴/ml) in 100 µl of DMEM high glucose media supplemented with L-glutamine (0.29 mg/ml), 10% heat-inactivated FBS, and 500 µg/ml each of geneticin, penicillin and streptomycin in 96-well U-bottom plates. We then incubated the plates at 37°C, 5% CO₂ overnight to allow cells to attach. We then decanted the media from the plates and replaced it with 100 µl fresh media per well. For testing, we diluted antibodies in assay media (DMEM high glucose media supplemented with L-glutamine, 10% heat-inactivated FBS, 200 µg/ml

BSA and 500 µg/ml each of geneticin, penicillin and streptomycin) to the desired final concentration and added 50 µl per well. All samples were performed in triplicate. We then incubated the plates at 37°C for ten minutes. We diluted [¹²⁵I]-IGF-I to a concentration of 1 µCi/ml in assay media and added 50 µl per well of the plate. As a control for background radioactivity, we added cold IGF-I to a final concentration of 100 ng/ml. We incubated the plates for 10 minutes at 37°C, decanted the media by blotting gently onto paper towels and washed twice with assay media. We then lysed the cells by adding 50 µl 0.1 N NaOH, 0.1% SDS and shaking the plates for five minutes at room temperature. We then transferred the samples to a scintillation plate, added 150 µl OptiPhase Supermix and read the signal on a Wallac Micro-Beta counter.

Table III and Figure 3 show the results of this experiment performed with three representative antibodies of the invention. This experiment demonstrated that antibodies of the invention specifically inhibit binding of [¹²⁵I]-IGF-I to cells overexpressing IGF-IR.

Table III

Monoclonal Antibody	IC ₅₀
2.12.1	0.45 µg/ml
2.13.2	0.18 µg/ml
4.9.2	0.1 µg/ml

EXAMPLE V: Epitope Mapping Studies

Having demonstrated that the antibodies of the invention recognize IGF-IR, we performed epitope mapping studies with several antibodies of the invention. We focused these experiments particularly on the 2.12.1, 2.13.2, 2.14.3, and 4.9.2 antibodies.

We conducted BIAcore competition studies to determine whether the antibodies of this invention bind to the same or distinct site on the IGF-IR molecule. We bound the extracellular domain (ECD) of IGF-IR to a BIAcore sensorchip as described above in Example II. We bound a first antibody of the invention to this sensorchip-bound IGF-IR under saturating conditions. We then measured the ability of subsequent secondary antibodies of the invention to compete with the primary

antibody for binding to IGF-IR. This technique enabled us to assign the antibodies of this invention to different binding groups.

We performed this experiment with antibodies 2.12.1, 2.13.2, 2.14.3, and 4.9.2. We observed that 2.13.2 and 4.9.2 compete for the same site on the extracellular domain of IGF-IR. The other antibodies, 2.12.1 and 2.14.3, bind to sites on IGF-IR that are different from both each other and from the site bound by 2.13.2 and 4.9.2.

EXAMPLE VI: Species Crossreactivity of the Antibodies of the Invention

In order to determine the species crossreactivity of the antibodies of the invention, we performed several experiments including immunoprecipitation, antibody-mediating blocking of IGF-I-induced receptor phosphorylation and FACS analysis.

To perform immunoprecipitation experiments, we plated cells in DMEM high glucose media supplemented with L-glutamine (0.29 mg/ml), 10% heat-inactivated FBS, and 500 µg/ml each of geneticin, penicillin and streptomycin to 50% confluence in T25 flasks. We then added 100 µl of an antibody of the invention in Hank's buffered saline solution (HBSS; Gibco BRL) at a concentration of 1 µg/ml. We incubated the plates for 30 minutes at 37°C in an incubator and then stimulated the cells with IGF-I at 100 ng/ml for 10 minutes at room temperature. We lysed the cells in RIPA buffer (Harlow and Lane, *supra*) and immunoprecipitated IGF-IR with 2 µg of pan-specific SC-713 anti-IGF-IR antibody (Santa Cruz) plus protein A agarose beads for 1 hour at 4°C. We pelleted the beads and wash three times with PBS/T (PBS + 0.1% Tween-20) and then boiled the beads in 40 µl Laemmli buffer containing 5% βME.

The samples prepared as described above were then analyzed by Western blot. We loaded 12 µl of each sample per lane on 4-10% gradient Novex™ gels run with 1X MES buffer (Novex™). Gels were run at 150V for 1 hour or at 200V for approximately 30 minutes. We then transferred the gel to a membrane in Novex™ transfer buffer with 10% methanol either overnight at 100mA or for 1-1.5 hours at 250mA. We then allowed the membrane to dry completely and blocked at room temperature with TBS (Tris-buffered saline pH 8.0) containing Superblock (Pierce Chemical Co.). We added the IGF-IR blotting antibody SC713 (Santa Cruz) to detect immunoprecipitated IGF-IR.

This experiment was performed with antibodies of the invention, particularly 2.12.1, 2.13.2, 4.17.3 and 4.9.2, on cells from a variety of animals. We found that antibodies 2.12.1, 2.13.2 and 4.9.2 were able to bind human, but not canine, guinea pig, rabbit or IGF-IR. Further, these antibodies were able to bind
5 COS7 and Rhesus IGF-IR, both derived from old world monkeys, but not IGF-IR from the marmoset, which is a new world monkey. These experiments indicate that the antibodies are highly specific.

*Antibody-mediated Blocking of
IGF-I/IGF-IR Binding in Non-human Primates*

10 Following our observation that the antibodies of the invention recognize IGF-IR from old world monkeys, we also tested their ability to block IGF-I/IGF-IR binding in cells derived from these old world monkeys. We plated cells in DMEM high glucose media supplemented with L-glutamine, 10% heat-inactivated FBS, and 500 µg/ml each of geneticin, penicillin and streptomycin to 50% confluence
15 in T25 flasks. We then added an antibody of the invention, or media without antibody as a control, and stimulated the cells with IGF-I at 100 ng/ml for 10 minutes at room temperature. After stimulation, we lysed the cells and immunoprecipitated IGF-IR with pan-specific IGF-IR antibody SC713 as described above. We then performed Western blot analysis as described above using HRP-PY54 antibody to detect
20 phosphorylated tyrosine in the activated IGF-IR.

We observed that antibodies of this invention, in particular 2.13.2 and 4.9.2 could block IGF-I-induced phosphorylation of IGF-IR in both COS7 and Rhesus cells. The IC₅₀ for the observed inhibition was 0.02 µg/ml and 0.005 µg/ml for COS7 and Rhesus IGF-IR, respectively.

25 *Determination of Cross-species Affinity
of Antibodies of the Invention*

We performed FACS analysis to determine the affinity of the antibodies of the invention for IGF-IR from other animals, particularly the old world monkeys described above. We incubated aliquots of human and monkey cells (5x10⁵)
30 for 1 hour on ice with increasing concentrations of biotinylated anti-IGF-IR antibodies of the invention or with a biotinylated anti-keyhole limpet hemocyanin (KLH) antibody (Abgenix) as a negative control. We then incubated the samples for 30 minutes on ice with streptavidin-conjugated RPE (phycoerythrin). We measured

binding by flow cytometry and analyzed the histograms of fluorescence intensity (FI2-H) versus cell number (Counts) using CellQuest software. We calculated binding (K_d) for each antibody from graphs of mean fluorescence intensity versus antibody concentration. In most experiments, we measured binding in cultured
5 human MCF-7 cells and either rhesus or cynomologous tissue culture cells. We controlled for depletion of the antibody by measuring binding over a range of cell concentrations.

We performed the aforementioned FACS analysis to test the ability of antibodies of the invention, particularly 2.13.2 and 4.9.2, to bind human, rhesus and
10 cynomologous cells. We observed a half maximal binding (K_d) of 0.1 $\mu\text{g/ml}$ for all cell lines tested.

EXAMPLE VII: IGF-I Receptor Downregulation

We performed blocking experiments essentially as described above in Example IV up to the addition of [^{125}I]-labeled IGF-I. At this point, we boiled the
15 cells in 40 μl Laemmli buffer containing 50% me. We then analyzed the samples by western blot analysis as described above in Example VI and probed the blots with both pan-specific IGF-IR antibody SC713 to quantify the levels of IGF-IR and HRP-PY54 antibody to monitor the levels of phosphorylated tyrosine in the activated IGF-IR.

As observed previously (Example III), we observed blockage of IGF-I-induced phosphorylation of IGF-IR following the treatment of cells with an antibody of this invention (Figure 4). Further, we observed that this blockage of IGF-I-induced phosphorylation was followed by downregulation of the IGF-IR in these cells. See, e.g., Fig. 4. IGF-IR levels were maximally reduced 16 hours after stimulation with
25 IGF-I in the presence of an antibody of the invention.

EXAMPLE VIII: Effects of the Antibodies of the Invention on IGF-IR *in vivo*

We determined whether the effects of the antibodies of the invention on IGF-IR as described in the previous examples would occur *in vivo*. We induced tumors in athymic mice according to published methods (V.A. Pollack et al.,
30 "Inhibition of epidermal growth factor receptor-associated tyrosine phosphorylation in human carcinomas with CP-358,774: Dynamics of receptor inhibition in situ and antitumor effects in athymic mice," J. Pharmacol. Exp. Ther. 291:739-748 (1999).

Briefly, we injected IGF-IR-transfected NIH-3T3 cells (5×10^6) subcutaneously into 3-4 week-old athymic (*nu/nu*) mice with 0.2 ml of Matrigel preparation. We then injected mice with an antibody of the invention intraperitoneally after established (i.e. approximately 400 mm³) tumors formed.

5 After 24 hours, we extracted the tumors, homogenized them and determined the level of IGF-IR. To determine IGF-IR levels, we diluted the SC-713 antibody in Blocking buffer to a final concentration of $\mu\text{g/ml}$ and added 100 μl to each well of a Reacti-Bind Goat anti-rabbit (GAR) coated plate (Pierce). We incubated the plates at room temperature for 1 hour with shaking and then washed the plates five
10 times with wash buffer. We then weighed tumor samples that had been prepared as described above and homogenized them in lysis buffer (1 ml/100 mg). We diluted 12.5 μl of tumor extract with lysis buffer to a final volume of 100 μl and added this to each well of a 96-well plate. We incubated the plates at room temperature with shaking for 1-2 hours and then washed the plates five times with Wash buffer. We
15 then added 100 μl HRP-PY54 or biotinylated anti-IGF-IR antibody in Blocking buffer to each well and incubated at room temperature with shaking for 30 minutes. We then washed the plates five times with wash buffer and developed the plates. We developed the plates probed with HRP-PY54 by adding 100 μl of the TMB microwell substrate per well and stopped color development with the addition 100 μl 0.9 M
20 H₂SO₄. We then quantitated the signal by shaking for 10 seconds and measuring OD_{450nm}. The signal was normalized to total protein. We developed plates probed with anti-IGF-IR antibody by adding 100 μl of streptavidin-HRP diluted in Blocking buffer to each well, incubating at room temperature with shaking for 30 minutes and then continuing as described for HRP-PY54.

25 We observed that intraperitoneal injection of an antibody of this invention, particularly 2.13.2 and 4.9.2, resulted in inhibition of IGF-IR activity as measured by a decrease of [both IGF-IR phosphotyrosine (phosphorylated IGF-IR) and] total IGF-IR protein (Figure 6). In addition, we also observed a decrease in IGF-IR phosphotyrosine (phosphorylated IGF-IR) (Figure 5). Without wishing to be bound
30 by any theory, the decreased levels of IGF-IR phosphotyrosine may be due to the decreased levels of IGF-IR protein *in vivo* after treatment with the antibody or may be due to a combination of decreased levels of IGF-IR protein and a decrease in tyrosine phosphorylation on the IGF-IR that is present due to blocking of activation by ligand

(e.g., IGF-I or IGF-II). Furthermore, this inhibition was responsive to the dose of antibody injected (Figure 6). These data demonstrate that the antibodies of the invention are able to target the IGF-IR *in vivo* in a manner analogous to what we observed *in vitro*.

5 EXAMPLE IX: Growth Inhibition (TGI) of 3T3/IGF-IR Cell Tumors

 We tested whether anti-IGF-IR antibodies of the invention would function to inhibit tumor growth. We induced tumors as described above (Example VIII) and when established, palpable tumors formed (i.e. 250 mm³, within 6-9 days), we treated the mice with a single, 0.20 ml dose of antibody by intraperitoneal
10 injection. We measured tumor size by Vernier calipers across two diameters every third day and calculated the volume using the formula (length x [width]²)/2 using methods established by Geran, et al., "Protocols for screening chemical agents and natural products against animal tumors and other biological systems," Cancer Chemother. Rep. 3:1-104.

15 When we performed this analysis with an antibody of the invention, we found that a single treatment with antibody 2.13.2 alone inhibited the growth of IGF-IR-transfected NIH-3T3 cell-induced tumors (Figure 7, left panel). Furthermore, in combination studies with a single dose of 7.5 mg/kg intravenously-supplied adriamycin, we observed that administration of a single dose of 2.13.2 enhanced the
20 effectiveness of adriamycin, a known inhibitor of tumor growth. The combination of adriamycin with an antibody of the invention, 2.13.2, demonstrated a growth delay of 7 days versus treatment with the antibody or adriamycin alone (Figure 7, right panel).

EXAMPLE X: Relationship of Antibody Levels to IGF-IR Downregulation

 Tumors were induced in nude mice as described in Example VIII. The
25 mice were then treated with 125 µg of 2.13.2 by intraperitoneal injection, as described in Example VIII. Tumors were extracted and IGF-IR levels were measured by ELISA as described in Example VIII. Figure 8 shows the serum 2.13.2 antibody levels and IGF-IR receptor levels over time. The experiment demonstrates that the IGF-IR is down-regulated by the antibody and that the degree of IGF-IR inhibition is dose
30 proportional to the serum concentration of the antibody.

EXAMPLE XI: Growth Inhibition of 3T3/IGF-IR Tumors with Multiple Dosing of Antibody in Combination with Adriamycin

Tumors were induced in nude mice as described in Example IX. Mice with established subcutaneous tumors of approximately 250 mm³ were treated on days 1, 8, 15 and 22 with various amounts of 2.13.2 antibody (i.p.) or 7.5 mg/kg adriamycin (i.v.), either as single agents or in combination, as described in Example IX. Figure 9 shows the tumor size in relation to the various treatments over time. The experiment demonstrates that treatment with an anti-IGF-IR antibody given once every seven days inhibits tumor cell growth and enhances inhibition of tumor cell growth in combination with adriamycin, a known tumor inhibitor.

EXAMPLE XII: Growth Inhibition of Large Tumors

Tumors were induced in nude mice as described in Example IX. Mice with large established subcutaneous tumors of slightly less than 2000 mm³ were treated on days 1 and 8 with various amounts of 2.13.2 antibody (i.p.) or 7.5 mg/kg adriamycin (i.v.), either as single agents or in combination, as described in Example IX. Figure 10 shows the tumor size in relation to the various treatments over time. Control, antibody alone and adriamycin alone animal groups were terminated at day 5, when the tumor size exceeded 2000 mm³. The experiment demonstrates that treatment with an anti-IGF-IR antibody in combination with adriamycin is highly efficacious against large tumors when multiple doses are given.

EXAMPLE XIII: Growth Inhibition of Colorectal Cell Tumors

Tumors were induced in nude mice as described in Example IX except that Colo 205 cells (ATCC CCL 222) were used. Colo 205 cells are human colorectal adenocarcinoma cells. Mice with established subcutaneous tumors of approximately 250 mm³ were treated with various amounts of 2.13.2 antibody (i.p.) or with 100 mg/kg 5-fluorodeoxyuridine (5-FU, i.v.), either as single agents or in combination, as described in Example IX. Figure 11 shows the tumor size in relation to the various treatments over time. The experiment demonstrates that treatment with an anti-IGF-IR antibody given once inhibits human colorectal cancer cell growth when provided as a single agent and enhances the effectiveness of 5-FU, a known tumor inhibitor.

Mice with established Colo 205 tumors were treated on days 1, 8, 15 and 22 with 500 µg 2.13.2 (i.p.), 100 mg/kg 5-FU (i.v.) or a combination thereof.

Figure 12 shows the tumor size in relation to the various treatments over time. The experiment demonstrates that treatment with an anti-IGF-IR antibody given once every seven days inhibits human colorectal cancer cell growth and enhances the effectiveness of 5-FU.

5 EXAMPLE XIV: Growth Inhibition of Breast Cancer Cell Tumors

Nude mice as described in Example VIII were implanted with biodegradable estrogen pellets (0.72 mg 17- β -estradiol/pellet, 60 day release; Innovative Research of America). After 48 hours, tumors were induced in nude mice essentially as described in Example IX except that MCF-7 cells (ATCC HTB-22)
10 were used. MCF-7 cells are estrogen-dependent human breast carcinoma cells. Mice with established subcutaneous tumors of approximately 250 mm³ were treated with 50 μ g 2.13.2 antibody (i.p.) on days 1, 4, 7, 10, 13, 16, 19 and 22 (q3dx7) or with 6.25 mg/kg taxol (i.p.) on days 1, 2, 3, 4, 5 (q1dx5), either as single agents or in combination, essentially as described in Example IX. Figure 13 shows the tumor size
15 in relation to the various treatments over time. The experiment demonstrates that treatment with an anti-IGF-IR antibody by itself inhibits human breast cancer cell growth when administered once every three days and also enhances the effectiveness of taxol, a known breast cancer inhibitor, when given in combination.

Mice having established tumors from MCF-7 cells as described
20 immediately above were treated on day 1 with various amounts of 2.13.2 antibody (i.p.) alone or with 3.75 mg/kg adriamycin (i.v.), essentially as described in Example IX. Figure 14 shows the tumor size in relation to the various treatments over time. The experiment demonstrates that a single treatment with an anti-IGF-IR antibody by itself inhibits human breast cancer cell growth and enhances the effectiveness of
25 adriamycin, a known tumor inhibitor.

Mice having established tumors from MCF-7 cells as described immediately above were treated with 250 μ g 2.13.2 antibody (i.p.) on days 1, 8, 15 and 23 or with a biodegradable tamoxifen pellet (25 mg/pellet, free base, 60 day release, Innovative Research of America), either as single agents or in combination,
30 essentially as described in Example IX. The tamoxifen pellet was implanted on day 1 after the tumor was established. Figure 15 shows the tumor size in relation to the various treatments over time. The experiment demonstrates that treatment with an

anti-IGF-IR antibody administered once every seven days inhibits human breast cancer growth by itself and enhances the effectiveness of tamoxifen, a known tumor inhibitor.

5 EXAMPLE XVI: Growth Inhibition of Epidermoid Carcinoma Cell Tumors

Tumors were induced in nude mice essentially as described in Example IX except that A431 cells (ATCC CRL 1555) were used. A431 cells are human epidermoid carcinoma cells that overexpress EGFR. Mice with established subcutaneous tumors of approximately 250 mm³ were treated on days 1, 8, 15, 22 and
10 29 with 500 µg 2.13.2 antibody (i.p.) or were treated once daily for 27 days with 10 mg/kg CP-358,774 given orally (p.o.), either as single agents or in combination, as described in Example IX. CP-358,774 is described in United States Patent 5,747,498 and Moyer et al., Cancer Research 57: 4838-4848 (1997), herein incorporated by reference. Figure 16 shows the tumor size in relation to the various treatments over
15 time. The experiment demonstrates that treatment with an anti-IGF-IR antibody enhances the effectiveness of CP-358,774, a known EGF-R tyrosine kinase inhibitor, for inhibiting the growth of a human epidermoid carcinoma tumor.

EXAMPLE XVII: Pharmacokinetics of Anti-IGF-IR Antibodies *in vivo*

To evaluate the pharmacokinetics of the anti-IGF-IR antibodies,
20 cynomolgus monkeys were injected intravenously with 3, 30 or 100 mg/kg of 2.13.2 antibody in an acetate buffer. Serum was collected from the monkeys at various time points and anti-IGF-IR antibody concentrations in the monkeys were determined for a period of up to ten weeks levels. To quantitate functional serum antibody levels, the extracellular domain of the human IGF-IR (IGF-I-sR, R&D Systems, Catalog #
25 391GR) was bound to 96-well plates. Monkey serum (diluted between 1:100 and 1:15,000) was added to the assay plates so that each sample would be within the linear range of the standard curve and incubated under conditions in which any anti-IGF-IR antibody would bind to IGF-I-sR. After washing the plates, a labeled anti-human IgG antibody was added to the plates and incubated under conditions in which the anti-
30 human IgG antibody would bind to the anti-IGF-IR antibody. The plates were then washed and developed, and a control standard curve and linear regression fits used to quantitate the amount of anti-IGF-IR antibodies. Figure 17 shows the concentration

of 2.13.2 in serum over time. The experiment demonstrates that the half-life of the anti-IGF-IR antibody is 4.6 to 7.7 days and has a volume distribution of 74-105 mL/kg. Further, the experiment demonstrates that the amounts given are dose-proportional in the monkey, which indicates that the anti-IGF-IR antibody has
5 saturated any available IGF-IR binding sites in the body even at the lowest dose of 3 mg/kg.

EXAMPLE XVIII: Combination Therapy of Anti-IGF-IR Antibody and Adriamycin
Downregulates IGF-IR *in vivo*

10 Tumors were induced in nude mice as described in Example IX. Mice with established subcutaneous tumors of approximately 400 mm³ were treated with a single injection of 250 µg 2.13.2 antibody (i.p.) or with 7.5 mg/kg adriamycin (i.v.), either as single agents or in combination, as described in Example IX. 72 hours after administration of the agents, tumors were extracted as described in Example VIII, and
15 equal amounts of the tumor extracts were subjected to sodium dodecyl phosphate polyacrylamide gel electrophoresis (SDS PAGE) and western blot analysis using the anti-IGF-IR antibody SC-713 (Santa Cruz). Figure 18 shows the amounts of IGF-IR in tumor cells in control animals (first three lanes of each panel), in animals treated with antibody alone (top panel), in animals treated with adriamycin alone (middle
20 panel) and in animals treated with antibody and adriamycin (lower panel). Each lane represents equal amounts of protein from individual tumors from individual mice. The experiment demonstrates that treatment with adriamycin alone has little effect on IGF-IR levels and that treatment with antibody alone shows some decrease in IGF-IR levels. Surprisingly, treatment with adriamycin and antibody together shows a
25 dramatic decrease in IGF-IR levels, demonstrating that adriamycin and antibody greatly downregulate IGF-IR levels.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although
30 the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

CLAIMS

We claim:

1. A humanized, chimeric or human monoclonal antibody or antigen-binding portion thereof that specifically binds to insulin-like growth factor I receptor (IGF-IR).

2. The antibody or portion thereof according to claim 1, wherein the IGF-IR is human.

10

3. The antibody or antigen-binding portion thereof according to any one of claims 1 or 2, wherein the antibody or portion thereof has at least one property selected from the group consisting of:

- a) does not bind to mouse, rat, dog or rabbit IGF-IR;
- 15 b) binds to cynomologous or rhesus IGF-IR but not to marmoset IGF-IR;
- c) inhibits the binding of IGF-I or IGF-II to IGF-IR.
- d) has a selectivity for IGF-IR that is at least 50 times greater than its selectivity for insulin receptor;
- 20 e) inhibits tumor growth *in vivo*;
- f) causes IGF-IR disappearance from the cell surface when incubated with a cell expressing IGF-IR;
- g) inhibits IGF-IR-induced tyrosine phosphorylation;
- h) binds to IGF-IR with a K_d of 8×10^{-9} M or less; and
- 25 i) has an off rate for IGF-IR of K_{off} of 10^{-4} or smaller.

4. The antibody or antigen-binding portion thereof according to claim 3, wherein the antibody or portion thereof has all of said properties.

30

5. The antibody or antigen-binding portion thereof according to any one of claims 1 or 2, wherein the antibody or portion thereof has at least one property selected from the group consisting of:

- a) cross-competes for binding to IGF-IR with an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- b) binds to the same epitope of IGF-IR as an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- 5 c) binds to the same antigen as that bound by the antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- d) binds to IGF-IR with substantially the same K_d as an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3; and
- 10 e) binds to IGF-IR with substantially the same off rate as an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3.
6. The antibody or antigen-binding portion thereof according to claim 5, wherein the antibody or portion thereof comprises all of said properties.
- 15 7. The antibody or antigen-binding portion thereof according to any one of claims 1-6, wherein said antibody or portion thereof inhibits binding between IGF-IR and IGF-I or IGF-II with an IC_{50} of less than 100 nM.
- 20 8. The antibody or antigen-binding portion thereof according to any one of claims 1-7, wherein said antibody or antigen-binding portion thereof comprises a variable region of a κ light chain, wherein the sequence of said variable region of said κ light chain comprises no more than ten amino acid changes from the amino acid sequence encoded by a germline V κ A30, A27 or O12 gene.
- 25 9. The antibody or antigen-binding portion thereof according to claim 8, wherein the variable region of the κ light chain comprises an amino acid sequence selected from the group consisting of the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 10, SEQ ID NO: 14, SEQ ID NO: 18 and SEQ ID NO: 22, or
- 30 an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions therefrom.
10. The antibody or antigen-binding portion thereof according to any one of claims 1-7, wherein said antibody or antigen-binding portion thereof comprises

a variable region of a heavy chain, wherein the sequence of said variable region comprises no more than eight amino acid changes from the amino acid sequence encoded by a germline V_H DP47, DP35, DP71 or VIV-4 gene.

5 11. The antibody or antigen-binding portion thereof according to claim 10, wherein the variable region of the heavy chain comprises an amino acid sequence selected from the group consisting of the amino acid sequence of SEQ ID NO: 4, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 20 and SEQ ID NO: 24, or an amino acid sequence having 1-10 amino acid insertions, deletions or
10 substitutions therefrom.

12. The antibody or antigen-binding portion thereof according to any one of claims 1-11 that is

 a) an immunoglobulin G (IgG), an IgM, an IgE, an IgA or an IgD
15 molecule, or is derived therefrom; or
 b) an Fab fragment, an F(ab')₂ fragment, an F_v fragment, a single chain antibody, a humanized antibody, a chimeric antibody or a bispecific antibody.

13. The antibody or antigen-binding portion thereof according to any
20 one of claims 1-12, wherein the antibody or portion thereof comprises an amino acid sequence of at least one CDR region from a variable region, wherein said variable region is selected from the group consisting of:

 a) a variable region of the light chain of an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
25 b) a variable region of a light chain comprising an amino acid sequence selected from SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 10, SEQ ID NO: 14, SEQ ID NO: 18 and SEQ ID NO: 22, or an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions therefrom;
 c) a variable region of the heavy chain of an antibody selected from
30 the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3; and
 d) a variable region of a heavy chain comprising an amino acid sequence selected from SEQ ID NO: 4, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID

NO:16, SEQ ID NO: 20 and SEQ ID NO: 24, or an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions therefrom.

14. The antibody or antigen-binding portion thereof according to
5 claim 13, wherein the antibody or portion thereof comprises all of the amino acid sequences of the CDR regions a variable region, wherein said variable region is selected from the group consisting of selected from the group consisting of:
- a) a variable region of the light chain of an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
 - 10 b) a variable region of a light chain comprising an amino acid sequence selected from SEQ ID NO: 2, SEQ ID NO: 6, SEQ ID NO: 10, SEQ ID NO: 14, SEQ ID NO: 18 and SEQ ID NO: 22, or an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions therefrom;
 - c) a variable region of the heavy chain of an antibody selected from
15 the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
 - d) a variable region of a heavy chain comprising an amino acid sequence selected from SEQ ID NO: 4, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO:16, SEQ ID NO: 20 and SEQ ID NO: 24, or an amino acid sequence having 1-10 amino acid insertions, deletions or substitutions; and
 - 20 e) the variable regions of the light chain and the heavy chain of an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3.

15. The antibody according to claim 1, wherein the antibody is
25 selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3.

16. The antibody according to claim 1, wherein said antibody comprises a heavy chain and a light chain, and wherein the amino acid sequences of the light chain and heavy chain are selected from the group consisting of:
- 30 a) the amino acid sequence of the heavy chain and the amino acid sequence of the light chain of 2.12.1;
 - b) the amino acid sequence of the heavy chain and the amino acid sequence of the light chain of 2.13.2;

c) the amino acid sequence of SEQ ID NO: 45 and the amino acid sequence of SEQ ID NO: 47; and

d) the amino acid sequence of SEQ ID NO: 49 and the amino acid sequence of SEQ ID NO: 51.

5

17. The antibody according to claim 1, wherein said antibody has an amino acid sequence comprising the amino acid sequences of the CDRs of antibodies 2.12.1 or 2.13.2, or CDRs of that antibody having no more than 5 conservative amino acid changes.

10

18. A pharmaceutical composition comprising the antibody or portion thereof according to any one of claims 1-17 and a pharmaceutically acceptable carrier.

19. The pharmaceutical composition according to claim 18, further comprising an antineoplastic, chemotherapeutic or anti-tumor agent.

15

20. A process for making an antibody that binds to IGF-IR, comprising the steps of:

a) immunizing a non-human mammal with an immunogen comprising IGF-IR, wherein the mammal is capable of expressing human antibodies in B cells of the animal;

20

b) isolating B cells from the mammal;

c) screening said B cells, or cell lines derived therefrom, to identify a cell line that produces antibodies that bind to IGF-IR;

25

d) culturing the cell line that expresses antibodies that bind to IGF-IR;

and

e) isolating antibodies that bind to IGF-IR from the cell line.

21. An isolated cell line that produces the antibody according to any one of claims 1-17.

30

22. The cell line according to claim 20 that produces an antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3, or wherein the antibody has the same amino acid sequences thereof.

- 5 23. A method of diagnosing the presence or location of an IGF-IR-expressing tumor in a subject in need thereof, comprising the steps of
- a) injecting the antibody according to any one of claims 1-17 into the subject,
 - b) determining the expression of IGF-IR in the subject by
 - 10 localizing where the antibody has bound,
 - c) comparing the expression in part (b) with that of a normal reference subject or standard, and
 - d) diagnosing the presence or location of the tumor.

- 15 24. A method of treating cancer in a human with an antibody or antigen-binding portion thereof that specifically binds to IGF-IR, comprising the step of administering to the human an amount of the antibody effective to treat said cancer.

- 20 25. A method of treating a patient in need thereof with the antibody or antigen-binding portion thereof according to any one of claims 1-17, comprising the step of administering to the patient an effective amount of the antibody.

- 25 26. The method according to either of claims 24 or 25, further comprising the step of administering an anti-neoplastic, anti-tumor, anti-angiogenic or chemotherapeutic agent.

- 30 27. An isolated nucleic acid molecule that comprises a nucleic acid sequence that encodes a heavy chain or antigen-binding portion thereof or a light chain or antigen-binding portion thereof of an antibody according to any one of claims 1-17.

28. The isolated nucleic acid molecule according to claim 27, wherein the nucleic acid molecule comprises a nucleic acid sequence selected from the group consisting of:

- 5 a) a nucleic acid sequence encoding at least one CDR region from the heavy chain of the antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- b) a nucleic acid sequence encoding the three CDR regions from the heavy chain of the antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- 10 c) a nucleic acid sequence encoding the amino acid sequence of the heavy chain or the antigen-binding portion thereof of the antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- d) a nucleic acid sequence encoding at least one CDR sequence from the light chain of the antibody selected from the group consisting of 2.12.1, 2.13.2, 15 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- e) a nucleic acid sequence encoding the three CDR sequences from the heavy chain of the antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- f) a nucleic acid sequence encoding the amino acid sequence of the 20 light chain or the antigen-binding portion thereof of the antibody selected from the group consisting of 2.12.1, 2.13.2, 2.14.3, 3.1.1, 4.9.2 and 4.17.3;
- g) a nucleic acid sequence encoding the amino acid sequence of selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20 and 22; and
- 25 h) a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21 and 23;

wherein said nucleic acid molecule optionally comprises a nucleic acid sequence encoding the amino acid sequence of SEQ ID NO: 28 or SEQ ID NO: 26.

- 30 29. A vector comprising the nucleic acid molecule according to either of claims 27 or 28, wherein the vector optionally comprises an expression control sequence operably linked to the nucleic acid molecule.

30. A host cell comprising the vector according to claim 29 or the nucleic acid molecule according to either of claims 27 or 28.

5 31. A method of making an anti-IGF-IR antibody or antigen-binding portion thereof, comprising culturing the host cell according to claim 30 or the cell line according to claim 21 under suitable conditions and recovering said antibody or antigen-binding portion.

10 32. A non-human transgenic animal comprising the nucleic acid according to either of claims 27 or 28, wherein the non-human transgenic animal expresses said nucleic acid.

15 33. A method of treating a subject in need thereof with an antibody or antigen-binding portion thereof that specifically binds to IGF-IR, comprising the steps of

(a) administering an effective amount of an isolated nucleic acid molecule encoding the heavy chain or the antigen-binding portion thereof, an isolated nucleic acid molecule encoding the light chain or the antigen-binding portion thereof, or both the nucleic acid molecules encoding the light chain and the heavy chain or
20 antigen-binding portions thereof; and

(b) expressing the nucleic acid molecule.

1/25

2.13.2K	GACATCCAGA	TGACCCAGTT	TCCATCCTCC	CTGTCTGCAT	CTGTAGGAGA	50
A30	GACATCCAGA	TGACCCAGTC	TCCATCCTCC	CTGTCTGCAT	CTGTAGGAGA	50
2.14.3K	-----	-----	----TCCTCC	CTGTCTGCAT	CTGTAGGAGA	26
2.12.1K	-----	-----	-----	-----TGCAT	CTGTAGGAGA	15
4.9.2K	GACATCCAGA	TGACCCAGTC	TCCATCCTCC	CTGTCTGCAT	CTGTAGGAGA	50
Consensus	GACATCCAGA	TGACCCAGTY	TCCATCCTCC	CTGTCTGCAT	CTGTAGGAGA	50
CDR1						
2.13.2K	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GGGCATTAGA	AAATGATTTAG	100
A30	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GGGCATTAGA	AAATGATTTAG	100
2.14.3K	CAGAGTCACC	TTCACCTGCC	GGGCAAGTCA	GGACATTAGA	CGTGATTTAG	76
2.12.1K	CAGAGTCACC	TTCACCTGCC	GGGCAAGTCA	GGACATTAGA	CGTGATTTAG	65
4.9.2K	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GGGCATTAGA	AAATGATTTAG	100
Consensus	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GGGCATTAGA	AAATGATTTAG	100
2.13.2K	GCTGGTATCA	GCAGAAACCA	GGGAAAGCCC	CTAAGCGCCT	GATCTATGCT	150
A30	GCTGGTATCA	GCAGAAACCA	GGGAAAGCCC	CTAAGCGCCT	GATCTATGCT	150
2.14.3K	GCTGGTATCA	GCAGAAACCA	GGGAAAGCIC	CTAAGCGCCT	GATCTATGCT	126
2.12.1K	GCTGGTATCA	GCAGAAACCA	GGGAAAGCIC	CTAAGCGCCT	GATCTATGCT	115
4.9.2K	GCTGGTATCA	GCAGAAACCA	GGGAAAGCCC	CTAAGCGCCT	GATCTATGCT	150
Consensus	GCTGGTATCA	GCAGAAACCA	GGGAAAGCVC	CTAAGCGCCT	GATCTATGCT	150
CDR2						
2.13.2K	GCATCCCGTT	TGCAAGAGGG	GGTCCCATCA	AGGTTTCAGCG	GCAGTGGATC	200
A30	GCATCCAGTT	TGCAAGAGGG	GGTCCCATCA	AGGTTTCAGCG	GCAGTGGATC	200
2.14.3K	GCATCCCGTT	TACAAAGTGG	GGTCCCATCA	AGGTTTCAGCG	GCAGTGGATC	176
2.12.1K	GCATCCCGTT	TACAAAGTGG	GGTCCCATCA	AGGTTTCAGCG	GCAGTGGATC	165
4.9.2K	GCATCCAAAT	TACACCGTGG	GGTCCCATCA	AGGTTTCAGCG	GCAGTGGATC	200
Consensus	GCATCCMRWT	TRCAMMGAGG	GGTCCCATCA	AGGTTTCAGCG	GCAGTGGATC	200
2.13.2K	TGGGACAGAA	TTCACTCTCA	CAATCAGCAG	CCTGCAGCCT	GAAGATTTTG	250
A30	TGGGACAGAA	TTCACTCTCA	CAATCAGCAG	CCTGCAGCCT	GAAGATTTTG	250
2.14.3K	TGGGACAGAA	TTCACTCTCA	CAATCAGCAG	CCTGCAGCCT	GAAGATTTTG	226
2.12.1K	TGGGACAGAA	TTCACTCTCA	CAATCAGCAG	CCTGCAGCCT	GAAGATTTTG	215
4.9.2K	TGGGACAGAA	TTCACTCTCA	CAATCAGCAG	CCTGCAGCCT	GAAGATTTTG	250
Consensus	TGGGACAGAA	TTCACTCTCA	CAATCAGCMG	CCTGCAGCCT	GAAGATTTTG	250
CDR3						
2.13.2K	CAACTTATTA	CTGTCTACAA	CATAATAGTT	ACCGTGCAG	TTTTCGGCCAG	300
A30	CAACTTATTA	CTGTCTACAG	CATAATAGTT	ACCC-TCCN-	-----	288
2.14.3K	CAACTTATTA	CTGTCTACAG	CATAATAATT	ATCCTCGGAC	GTTCGGCCAA	276
2.12.1K	CAACTTATTA	CTGTCTACAG	CATAATAATT	ATCCTCGGAC	GTTCGGCCAA	265
4.9.2K	CAACTTATTA	CTGTCTACAG	CATAATAGTT	ACCTCTCAC	TTTCGGCGGA	300
Consensus	CAACTTATTA	CTGTCTACAR	CATAATAATT	AYCCKYBSNS	KTTYGGCSRR	300
2.13.2K	GGGACCAAGC	TGGAGATCAA	AC----			322
A30	-----	-----	-----			288
2.14.3K	GGGACCGAGG	TGGAAATCAT	ACGAAC			302
2.12.1K	GGGACCGAGG	TGGAAATCAT	ACGAAC			291
4.9.2K	GGGACCAAGG	TGGAGATCAA	AC----			322
Consensus	GGGACCRAGS	TGGARATCAW	ACGAAC			326

FIG. 1A

2/25

4.17.3K	-----	-----	-----	-----	-----	AGGAGA	7
O12	GACATCCAGA	TGACCCAGTC	TCCATCCTCC	CTGTCTGCAT	CTGTAGGAGA		50
Consensus	GACATCCAGA	TGACCCAGTC	TCCATCCTCC	CTGTCTGCAT	CTGYAGGAGA		50
CDR1							
4.17.3K	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GAGCATTAGT	ACCTTTTAA		57
O12	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GAGCATTAGC	ACCTTTTAA		100
Consensus	CAGAGTCACC	ATCACTTGCC	GGGCAAGTCA	GAGCATTAGY	ASCTTTTAA		100
4.17.3K	ATTGGTATCA	GCAGAAACCA	GGGAAAGCCC	CTAAACTCCT	GATCCATGTT		107
O12	ATTGGTATCA	GCAGAAACCA	GGGAAAGCCC	CTAAGCTCCT	GATCCTATGCT		150
Consensus	ATTGGTATCA	GCAGAAACCA	GGGAAAGCCC	CTAARCTCCT	GATCCTATGYT		150
CDR2							
4.17.3K	GCATCCAGTT	TCAAGGTGG	GGTCCCATCA	AGGTTCAAGT	GCAGTGGATC		157
O12	GCATCCAGTT	TCAAGGTGG	GGTCCCATCA	AGGTTCAAGT	GCAGTGGATC		200
Consensus	GCATCCAGTT	TCAAGGTGG	GGTCCCATCA	AGGTTCAAGT	GCAGTGGATC		200
4.17.3K	TGGGACAGAT	TTCACTCTCA	CCATCAGCAG	TCTGCAACCT	GAAGATTTTG		207
O12	TGGGACAGAT	TTCACTCTCA	CCATCAGCAG	TCTGCAACCT	GAAGATTTTG		250
Consensus	TGGGACAGAT	TTCACTCTCA	CCATCAGCAG	TCTGCAACCT	GAAGATTTTG		250
CDR3							
4.17.3K	CAACTTACTA	CTGTCAACAG	AGTTACATG	CCCCACTCAC	TTTCGGCGGA		257
O12	CAACTTACTA	CTGTCAACAG	AGTTACAGTA	CCCC-TCH-	-----		288
Consensus	CAACTTACTA	CTGTCAACAG	AGTTACATR	CCCCAYYCH	TTTCGGCGGA		300
4.17.3K	GGGACCAAGG	TGGAGATCAA	AC				279
O12	-----	-----	--				288
Consensus	GGGACCAAGG	TGGAGATCAA	AC				322

FIG. 1B

3/25

6.1.1K	-----					
A27	GAAATTGTGT	TGACGCAGTC	TCCAGGCACC	CTGTCTTTGT	CTCCAGGGGA	50
Consensus	GAAATTGTGT	TGACGCAGTC	TCCAGGCACC	CTGTCTTTGT	CTCCAGGGGA	50
	CDR1					
6.1.1K	-AGAGCCACC	CTCTCCTGTA	GGGCCAGTCA	GAGTGTTGCG	GGCAGGTACT	49
A27	AAGAGCCACC	CTCTCCTGCA	GGGCCAGTCA	GAGTGTTAGC	AGCAGGTACT	100
Consensus	AAGAGCCACC	CTCTCCTGTA	GGGCCAGTCA	GAGTGTTGCG	GGCAGGTACT	100
6.1.1K	TAGCCTGGTA	CCAGCAGAAA	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	99
A27	TAGCCTGGTA	CCAGCAGAAA	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	150
Consensus	TAGCCTGGTA	CCAGCAGAAA	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	150
	CDR2					
6.1.1K	GGTGCATCCA	GCAGGGCCAC	TGGCATCCCA	GACAGGTTCA	GTGGCAGTGG	149
A27	GGTGCATCCA	GCAGGGCCAC	TGGCATCCCA	GACAGGTTCA	GTGGCAGTGG	200
Consensus	GGTGCATCCA	GCAGGGCCAC	TGGCATCCCA	GACAGGTTCA	GTGGCAGTGG	200
6.1.1K	GTCTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	CCTGAAGATT	199
A27	GTCTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	CCTGAAGATT	250
Consensus	GTCTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	CCTGAAGATT	250
	CDR3					
6.1.1K	TTGCAGTGTT	TTACTGTCAG	CAGTATGGTA	GTTCACCTCG	NACGTTCCGC	249
A27	TTGCAGTGTA	TTACTGTCAG	CAGTATGGTA	GTTCACCTCC	-----	290
Consensus	TTGCAGTGTT	TTACTGTCAG	CAGTATGGTA	GTTCACCTCG	NACGTTCCGC	300
6.1.1K	CAAGGGACCA	AGGTGGAAAT	CAAAC			274
A27	-----	-----	-----			290
Consensus	CAAGGGACCA	AGGTGGAAAT	CAAAC			325

FIG. 1C

4/25

2.12.1H	-----	-----	---GGGAGGC	TTGGTCAAGC	CTGGA	GGTC	26
DP35	CAGGTGCAGC	TGGTGGAGTC	TGGGGGAGGC	TTGGTCAAGC	CTGGAG	GGTC	50
Consensus	CAGGTGCAGC	TGGTGGAGTC	TGGGGGAGGC	TTGGTCAAGC	CTGGAG	GGTC	50
							CDR1
2.12.1H	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CAC	TTTCAGT	GACTACTATA	76
DP35	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CAC	TTTCAGT	GACTACTACA	100
Consensus	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CAC	TTTCAGT	GACTACTATA	100
2.12.1H	TGAGCTGGAT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAATG	GGTTTCATAC		126
DP35	TGAGCTGGAT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAATG	GGTTTCATAC		150
Consensus	TGAGCTGGAT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAATG	GGTTTCATAC		150
							CDR2
2.12.1H	ATTAGTAGTA	GTGGTAGTAC	CAC	AACTAC	GCAGACTCTG	TGAAGGGCCG	176
DP35	ATTAGTAGTA	GTGGTAGTAC	CAA	TAACTAC	GCAGACTCTG	TGAAGGGCCG	200
Consensus	ATTAGTAGTA	GTGGTAGTAC	CAA	TAACTAC	GCAGACTCTG	TGAAGGGCCG	200
2.12.1H	ATTACCATC	TCCAGGGACA	ACGCCAAGAA	CTCACTGTAT	CTGCAAATGA		226
DP35	ATTACCATC	TCCAGGGACA	ACGCCAAGAA	CTCACTGTAT	CTGCAAATGA		250
Consensus	ATTACCATC	TCCAGGGACA	ACGCCAAGAA	CTCACTGTAT	CTGCAAATGA		250
2.12.1H	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTGT	ATTACTGTGT	GAGAGATGGA		276
DP35	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTGT	ATTACTGTGC	GAGAGATGGA		296
Consensus	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTGT	ATTACTGTGT	GAGAGATGGA		300
							CDR3
2.12.1H	GTGGAAACTA	CTTTTACTA	CTACTACTAC	GGTATGGACG	TCTGGGGCCA		326
DP35	-----	-----	-----	-----	-----		296
Consensus	GTGGAAACTA	CTTTTACTA	CTACTACTAC	GGTATGGACG	TCTGGGGCCA		350
2.12.1H	AGGGACCACG	GTCACCGTCT	CCTCAG				352
DP35	-----	-----	-----				296
Consensus	AGGGACCACG	GTCACCGTCT	CCTCAG				376

FIG. 2A

5/25

PF2-2.14.3H.DNA	-----	GGGCCCAGGA CTGGTGAAGC CTTCCGAGAC	30
VIV-4/4.35	CAGGTGCAGC TGCAGGAGTC	GGGCCCAGGA CTGGTGAAGC CTTCCGAGAC	50
Consensus	CAGGTGCAGC TGCAGGAGTC	GGGCCCAGGA CTGGTGAAGC CTTCCGAGAC	50
CDR1			
PF2-2.14.3H.DNA	CCTGTCCCTC ACCTGCACTG TCTCTGGTGG CTCCATCAGT	ATTACTACT	80
VIV-4/4.35	CCTGTCCCTC ACCTGCACTG TCTCTGGTGG CTCCATCAGT	ATTACTACT	100
Consensus	CCTGTCCCTC ACCTGCACTG TCTCTGGTGG CTCCATCAGT	ATTACTACT	100
CDR1			
PF2-2.14.3H.DNA	GGAGCTGGAT CCGGCAGCCC GCCGGAAGG GACTGGAGTG GATTGGGCGT		130
VIV-4/4.35	GGAGCTGGAT CCGGCAGCCC GCCGGAAGG GACTGGAGTG GATTGGGCGT		150
Consensus	GGAGCTGGAT CCGGCAGCCC GCCGGAAGG GACTGGAGTG GATTGGGCGT		150
CDR2			
PF2-2.14.3H.DNA	ATCTATACCA GTGGGAGCC CAACTACAAC CCCTCCCTCA AGAGTCGAGT		180
VIV-4/4.35	ATCTATACCA GTGGGAGCC CAACTACAAC CCCTCCCTCA AGAGTCGAGT		200
Consensus	ATCTATACCA GTGGGAGCC CAACTACAAC CCCTCCCTCA AGAGTCGAGT		200
PF2-2.14.3H.DNA	CACCATGTCA GTAGACACGT CCAAGAACCA GTTCTCCCTG AAGCTGACT		230
VIV-4/4.35	CACCATGTCA GTAGACACGT CCAAGAACCA GTTCTCCCTG AAGCTGACT		250
Consensus	CACCATGTCA GTAGACACGT CCAAGAACCA GTTCTCCCTG AAGCTGACT		250
PF2-2.14.3H.DNA	CTGTGACCGC CGCGGACACG GCCGTGTATT ACTGTGCGT	AACGATTTT	280
VIV-4/4.35	CTGTGACCGC CGCGGACACG GCCGTGTATT ACTGTGCGT	-----	288
Consensus	CTGTGACCGC CGCGGACACG GCCGTGTATT ACTGTGCGT	AACGATTTT	300
CDR3			
PF2-2.14.3H.DNA	GGAGTGGTTA TTATCTTTGA CTACTGGGGC CAGCGAACCC TGGTCACCGT		330
VIV-4/4.35	-----	AGAGAB----	294
Consensus	GGAGTGGTTA TTATCTTTGA CTACTGGGGC CAGCGAACCC TGGTCACCGT		350
PF2-2.14.3H.DNA	CTCCTCAG		338
VIV-4/4.35	-----		294
Consensus	CTCCTCAG		358

FIG. 2B

6/25

6.1.1H	GAGGTGCAGC	TGTTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGTC	50
4.9.2H	GAGGTGCAGC	TGTTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGTC	50
DP47	GAGGTGCAGC	TGTTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGTC	50
2.13.2H	GAGGTGCAGC	TGTTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGTC	50
Consensus	GAGGTGCAGC	TGTTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGTC	50
CDR1						
6.1.1H	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CACCTTTAGC	AGCTATGCCA	100
4.9.2H	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CACCTTTAGC	AGCTATGCCA	100
DP47	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CACCTTTAGC	AGCTATGCCA	100
2.13.2H	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CACCTTTAGC	AGCTATGCCA	100
Consensus	CCTGAGACTC	TCCTGTGCAG	CCTCTGGATT	CACCTTTAGC	AGCTATGCCA	100
CDR1						
CDR2						
6.1.1H	TGAGCTGGGT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAGTG	GGTCTCAGCT	150
4.9.2H	TGAGCTGGGT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAGTG	GGTCTCAGCT	150
DP47	TGAGCTGGGT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAGTG	GGTCTCAGCT	150
2.13.2H	TGAGCTGGGT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAGTG	GGTCTCAGCT	150
Consensus	TGAGCTGGGT	CCGCCAGGCT	CCAGGGAAGG	GGCTGGAGTG	GGTCTCAGCT	150
CDR2						
6.1.1H	ATTAGTGGTA	GTGGTGGTAG	TACATACTAC	GCAGACTCCG	TGAAGGGCCG	200
4.9.2H	ATTAGTGGTA	GTGGTGGTAG	TACATACTAC	GCAGACTCCG	TGAAGGGCCG	200
DP47	ATTAGTGGTA	GTGGTGGTAG	TACATACTAC	GCAGACTCCG	TGAAGGGCCG	200
2.13.2H	ATTAGTGGTA	GTGGTGGTAG	TACATACTAC	GCAGACTCCG	TGAAGGGCCG	200
Consensus	ATTAGTGGTA	GTGGTGGTAG	TACATACTAC	GCAGACTCCG	TGAAGGGCCG	200
CDR3						
6.1.1H	G TTCACCATC	TCCAGAGACA	ATTCCAGGAA	CACGCTGTAT	CTGCAAATGA	250
4.9.2H	G TTCACCATC	TCCAGAGACA	ATTCCAGGAA	CACGCTGTAT	CTGCAAATGA	250
DP47	G TTCACCATC	TCCAGAGACA	ATTCCAGGAA	CACGCTGTAT	CTGCAAATGA	250
2.13.2H	G TTCACCATC	TCCAGAGACA	ATTCCAGGAC	CACGCTGTAT	CTGCAAATGA	250
Consensus	G TTCACCATC	TCCAGAGACA	ATTCCAGGAA	CACGCTGTAT	CTGCAAATGA	250
CDR3						
6.1.1H	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTAT	ATTACTGTGC	GAAAGATC--	298
4.9.2H	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTAT	ATTACTGTGC	GAAAGATCTG	300
DP47	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTAT	ATTACTGTGC	GAAAGATCTG	296
2.13.2H	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTAT	ATTACTGTGC	GAAAGATCTT	300
Consensus	ACAGCCTGAG	AGCCGAGGAC	ACGGCCGTAT	ATTACTGTGC	GAAAGATCTK	300
CDR3-for 4.9.2 and 2.13.2						
6.1.1H	-----	-----	-----	-----	-----C-	299
4.9.2H	GGCTACGGTG	ACTTTTACTA	CTACTACTAC	GGTATGGACG	TCTGGGGCCA	350
DP47	-----	-----	-----	-----	-----	296
2.13.2H	GGCTGGTCCG	ACTCTTACTA	CTACTACTAC	GGTATGGACG	TCTGGGGCCA	350
Consensus	GGCTRSKSYG	ACTYTTACTA	CTACTACTAC	GGTATGGACG	TCTGGGGCCA	350
CDR3-for 6.1.1						
6.1.1H	AGGGACTACG	GTGATTATGA	GTTGGTTCGA	CCCCTGGGGC	CAGGGAACCC	349
4.9.2H	AGGGACCAC-	-----	-----	-----	-----	359
DP47	-----	-----	-----	-----	-----	296
2.13.2H	AGGGACCAC-	-----	-----	-----	-----	359
Consensus	AGGGACYACG	GTGATTATGA	GTTGGTTCGA	CCCCTGGGGC	CAGGGAACCC	400

FIG. 2C-1

7/25


6.1.1H	TGGTCACCGT CTCCTCAG	367
4.9.2H	-GGTCACCGT CTCCTCAG	376
DP47	-----	296
2.13.2H	-GGTCACCGT CTCCTCAG	376
Consensus	TGGTCACCGT CTCCTCAG	418

FIG. 2C-2

4.17.3H	-----	CCCAGGA CTGGTGAAGC CTTCGGAGAC	27
DP71	CAGGTGCAGC TGCAGGAGTC GGGCCCAGGA CTGGTGAAGC CTTCGGAGAC		50
Consensus	CAGGTGCAGC TGCAGGAGTC GGGCCCAGGA CTGGTGAAGC CTTCGGAGAC		50
		CDR1	
4.17.3H	CCTGTCCCTC ACCTGCACTG TCTCTGGTGG CTCCATCAGT AGTTACTACT		77
DP71	CCTGTCCCTC ACCTGCACTG TCTCTGGTGG CTCCATCAGT AGTTACTACT		100
Consensus	CCTGTCCCTC ACCTGCACTG TCTCTGGTGG CTCCATCAGT AGTTACTACT		100
		CDR1	
4.17.3H	GGAGCTGGAT CCGGCAGCCC CCAGGGAAGG GACTGGAGTG GATTGGGTAT		127
DP71	GGAGCTGGAT CCGGCAGCCC CCAGGGAAGG GACTGGAGTG GATTGGGTAT		150
Consensus	GGAGCTGGAT CCGGCAGCCC CCAGGGAAGG GACTGGAGTG GATTGGGTAT		150
		CDR2	
4.17.3H	ATCTATTACA GTGGGAGCAC CAACTACAAC CCCTCCCTCA AGAGTCGAGT		177
DP71	ATCTATTACA GTGGGAGCAC CAACTACAAC CCCTCCCTCA AGAGTCGAGT		200
Consensus	ATCTATTACA GTGGGAGCAC CAACTACAAC CCCTCCCTCA AGAGTCGAGT		200
4.17.3H	CACCATATCA GTAGACACGT CCAAGAACCA GTTCTCCCTG AAGCTGAGTT		227
DP71	CACCATATCA GTAGACACGT CCAAGAACCA GTTCTCCCTG AAGCTGAGCT		250
Consensus	CACCATATCA GTAGACACGT CCAAGAACCA GTTCTCCCTG AAGCTGAGTT		250
		CDR3	
4.17.3H	CTGTGACCGC TGCGGACACG GCCGTGTATT ACTGTGCCAG GACGTATAGC		277
DP71	CTGTGACCGC TGCGGACACG GCCGTGTATT ACTGTGCC--- GA-----		289
Consensus	CTGTGACCGC TGCGGACACG GCCGTGTATT ACTGTGCCAG GACGTATAGC		300
4.17.3H	AGTTCGTTCT ACTACTACGG TATGGACGTC TGGGGCCAAG GGACCACGGT		327
DP71	----- GA----- GA-----		293
Consensus	AGTTCGTTCT ACTACTACGG TATGGACGTC TGGGGCCAAG GGACCACGGT		350
4.17.3H	CACCGTCTCC TCAG		341
DP71	-----		293
Consensus	CACCGTCTCC TCAG		364

FIG. 2D

8/25



9/25




FIG. 4

10/25




FIG. 5




FIG. 6

11/25




FIG. 7

12/25


Treatment: 125 μ g dose

FIG. 8

13/25




FIG. 9

14/25




FIG. 10

15/25




FIG. 11

16/25




FIG. 12

17/25




FIG. 13

Combination




FIG. 14

19/25




FIG. 15

20/25




FIG. 16

21/25




FIG. 17

22/25




FIG. 18

23/ 25

Clone	C domain mutations	FR mutation	CDR mutation	Change in Cys	Change in glycosylation
2.13.2 Heavy	0	3	8	0	0
2.13.2 Light	0	1	4	1 (CDR3)	0
2.12.2 Heavy	0	2	8	0	0
2.12.2 Light	0	3	5	0	0

FIG. 19A

PF2 2.13.2 Heavy chain (DP-47 (3-23)/D6-19/JH6) +

MEFGLSWLFL VAILKGVCCE VQLLESGGGL VQPGGSLRLS CTASGFTFSS YAMNWRQAP GKGLEWVSAI SGSGGTFYA DSVKGRFTIS RDNSRTLLYL ++

MEFGLSWLFL VAILKGVCCE VQLLESGGGL VQPGGSLRLS CAASGFTFSS YAMNWRQAP GKGLEWVSAI SGSGGTFYA DSVKGRFTIS RDNSKNTLLYL

QMNSLRAEDT AVTYCAK--D LGWSDSYIYY YGMDVWGQGT TTVSSASTK GPSVFPLAPC SRSTSESTAA LGCLVKDYFP EPVTVSWNSG ALTSGVHTFP

QMNSLRAEDT AVTYCAKGIS SGW--YYYY YGMDVWGQGT TTVSSASTK GPSVFPLAPC SRSTSESTAA LGCLVKDYFP EPVTVSWNSG ALTSGVHTFP

AVLQSSGLYS LSSVVTVPSS NFGTQTYTCN VDHKPSNTKV DKTVERKCCV ECPPCPAPPV AGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVQF

AVLQSSGLYS LSSVVTVPSS NFGTQTYTCN VDHKPSNTKV DKTVERKCCV ECPPCPAPPV AGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVQF

NWYVDGVEVH NAKTKPREEQ FNSTFRVSV LTVVHQDWLN GKEYKCKVSN KGLPAPIEKT ISKTKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS

NWYVDGVEVH NAKTKPREEQ FNSTFRVSV LTVVHQDWLN GKEYKCKVSN KGLPAPIEKT ISKTKGQPRE PQVYTLPPSR EEMTKNQVSL TCLVKGFYPS

DIAVEESNG QPENNYKTTT PMLDSDGSFF LYSKLTVDKS RWQGNVFSC SVMHEALHNH YTKSLSLSP GK

DIAVEESNG QPENNYKTTT PMLDSDGSFF LYSKLTVDKS RWQGNVFSC SVMHEALHNH YTKSLSLSP GK

FIG. 19B

PF2 2.13.2 LC (A30/JK2) + * *

MDMRVPAQLL GLLLLWPPGA RCDIQMTQFP SSLSASVGDR VTITCRASQG IRNDLGWYQQ KPGKAPKRLI YAAASRLHRGV PSRPSGSGSG TEFTLTISSL

MDMRVPAQLL GLLLLWPPGA RCDIQMTQSP SSLSASVGDR VTITCRASQG IRNDLGWYQQ KPGKAPKRLI YAAASLQSGV PSRPSGSGSG TEFTLTISSL

QPEDFATYYC LQNSYPCSP GQGTKLEIKR TVAAPSVFIF PPSDEQLKSG TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYSLSST

QPEDFATYYC LQNSYPYTF GQGTKLEIKR TVAAPSVFIF PPSDEQLKSG TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYSLSST

LTLSKADYEK HKVYACEVTH QGLSSPVTKS FNRGEC

LTLSKADYEK HKVYACEVTH QGLSSPVTKS FNRGEC

FIG. 19C

PF2 2.12.1 Heavy chain (DP-35 (3-11)/D3-3/JH6) + **

MEFGLSWVPL VALIKGVQCQ AQLVESGGGL VKPGGSLRLS CAASGFTFSD YMSWIRQAP GKGLEWVSYI SSGSTRDYA DSVKGRFTIS RDNAKNSLYL

MEFGLSWVPL VALIKGVQCQ VQLVESGGGL VKPGGSLRLS CAASGFTFSD YMSWIRQAP GKGLEWVSYI SSGSTIYYA DSVKGRFTIS RDNAKNSLYL

QMSLRAEDT AVTYCVR--D GVETTF-YYY YYGMDVWGQG TTVTVSSAST KGPSVFFLAP CSRSTSESTA ALGCLVKDYF PEPVTVSWNS GALTSGVHTF

QMSLRAEDT AVTYCARVLR FLEWLLYYY YYGMDVWGQG TTVTVSSAST KGPSVFFLAP CSRSTSESTA ALGCLVKDYF PEPVTVSWNS GALTSGVHTF

PAVLQSSGLY SLSSVTVTPS SNFGTQTYTC NVDHKPSNTK VDKTVERKCC VECPPCPAPP VAGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVQ

PAVLQSSGLY SLSSVTVTPS SNFGTQTYTC NVDHKPSNTK VDKTVERKCC VECPPCPAPP VAGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVQ

FNWYVDGVEV HNAKTKPREE QFNSTFRVVS VLTVVHQDWL NGKEYKCKVS NKGLPAPIEK TISKTKGQPRE PQVYTLPPS REEMTKNQVS LTCLVKGFYP

FNWYVDGVEV HNAKTKPREE QFNSTFRVVS VLTVVHQDWL NGKEYKCKVS NKGLPAPIEK TISKTKGQPRE PQVYTLPPS REEMTKNQVS LTCLVKGFYP

SDIAVEWESN GOPENNYKTT PPMLDSGSGF FLYSKLTVDK SRWQQGNVFS CSVMHEALHN HYTKSLSLSP GK

SDIAVEWESN GOPENNYKTT PPMLDSGSGF FLYSKLTVDK SRWQQGNVFS CSVMHEALHN HYTKSLSLSP GK

FIG. 19D

25/25

PF2.12.1 Light chain (A30/Jk1)
 NDMRVPAQLL GLLLLWPPGA RCDIQMTQSP SLSASVGDR VTFTCRASQD IRNDLGWYQQ KPGKAPKRLI YAA SRLQSGV PSRFSGSGSG TEFTLTSSSL
 NDMRVPAQLL GLLLLWPPGA RCDIQMTQSP SLSASVGDR VTITCRASQ IRNDLGWYQQ KPGKAPKRLI YAASSLQSGV PSRFSGSGSG TEFTLTSSSL
 QPEDFATYYC LQHNNYPRTF GQGTEVEIIR TVAAPS VFIF PPSDEQLKSG TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYSLSSST
 QPEDFATYYC LQHNSYPWT GQGTEVEIIR TVAAPS VFIF PPSDEQLKSG TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYSLSSST
 LTLSKADYEK HKVYACEVTH QGLSSPVTKS FNRGEC
 LTLSKADYEK HKVYACEVTH QGLSSPVTKS FNRGEC

FIG. 19E

SEQUENCE LISTING

<110> ABGENIX, INC.
PFIZER, INC.

<120> ANTIBODIES TO INSULIN-LIKE GROWTH FACTOR I RECEPTOR

<130> ABX-PF2

<140>
<141>

<150> 60/259,927
<151> 2001-01-05

<160> 60

<170> PatentIn Ver. 2.1

<210> 1
<211> 291
<212> DNA
<213> Homo sapiens

<400> 1
tgcacatctgta ggagacagag tcaccttcac ttgccgggca agtcaggaca ttagacgtga 60
tttaggctgg tatcagcaga aaccagggaa agtccttaag cgcctgatct atgctgcatc 120
ccgtttacaa agtgggggtcc catcaagggt cagcggcagt ggatctggga cagaattcac 180
tctcacaatc agcagcctgc agcctgaaga ttttgcaact tattactgtc tacagcataa 240
taattatcct cggacgttcg gccaaaggac cgaggtggaa atcatacgaa c 291

<210> 2
<211> 136
<212> PRT
<213> Homo sapiens

<400> 2
Ala Ser Val Gly Asp Arg Val Thr Phe Thr Cys Arg Ala Ser Gln Asp
1 5 10 15
Ile Arg Arg Asp Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro
20 25 30
Lys Arg Leu Ile Tyr Ala Ala Ser Arg Leu Gln Ser Gly Val Pro Ser
35 40 45
Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser
50 55 60
Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn
65 70 75 80
Asn Tyr Pro Arg Thr Phe Gly Gln Gly Thr Glu Val Glu Ile Ile Arg
85 90 95
Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
100 105 110

2/31

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 115 120 125

Pro Arg Glu Ala Lys Val Gln Trp
 130 135

<210> 3
 <211> 352
 <212> DNA
 <213> Homo sapiens

<400> 3
 gggaggcttg gtcaagcctg gaggtccctg agactctcct gtgcagcctc tggattcact 60
 ttcagtgact actatatgag ctggatccgc caggctccag ggaaggggct ggaatgggtt 120
 tcatacatta gtagtagtgg tagtaccaga gactacgcag actctgtgaa gggccgattc 180
 accatctcca gggacaacgc caagaactca ctgtatctgc aaatgaacag cctgagagcc 240
 gaggacacgg ccgtgtatta ctgtgtgaga gatggagtgg aaactacttt ttactactac 300
 tactacggta tggacgtctg gggccaaggg accacggtca ccgtctcctc ag 352

<210> 4
 <211> 174
 <212> PRT
 <213> Homo sapiens

<400> 4
 Gly Arg Leu Gly Gln Ala Trp Arg Ser Leu Arg Leu Ser Cys Ala Ala
 1 5 10 15
 Ser Gly Phe Thr Phe Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala
 20 25 30
 Pro Gly Lys Gly Leu Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser
 35 40 45
 Thr Arg Asp Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg
 50 55 60
 Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala
 65 70 75 80
 Glu Asp Thr Ala Val Tyr Tyr Cys Val Arg Asp Gly Val Glu Thr Thr
 85 90 95
 Phe Tyr Tyr Tyr Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr
 100 105 110
 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160

3/31

Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ser Cys Ala
 165 170

<210> 5
 <211> 322
 <212> DNA
 <213> Homo sapiens

<400> 5
 gacatccaga tgacccagtt tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60
 atcacttgcc gggcaagtca gggcattaga aatgatttag gctgggtatca gcagaaacca 120
 gggaaagccc ctaagcgctt gatctatgct gcatcccggt tgcacagagg ggtcccatca 180
 aggttcagcg gcagtggatc tgggacagaa ttcactctca caatcagcag cctgcagcct 240
 gaagattttg caacttatta ctgtttacaa cataatagtt acccgtgcag ttttgccag 300
 gggaccaagc tggagatcaa ac 322

<210> 6
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 6
 Asp Ile Gln Met Thr Gln Phe Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Asp
 20 25 30
 Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
 35 40 45
 Tyr Ala Ala Ser Arg Leu His Arg Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Tyr Pro Cys
 85 90 95
 Ser Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 7
 <211> 375
 <212> DNA
 <213> Homo sapiens

<400> 7
 aggtgcagct gttggagtct gggggaggct tggtagagcc tgggggggtcc ctgagactct 60
 cctgtacagc ctctggattc accttttagca gctatgccat gaactggggtc cgccaggctc 120
 caggggaagg gctggagtgg gtctcagcta ttagtggttag tggtaggtacc acattctacg 180
 cagactccgt gaagggccgg ttcaccatct ccagagacaa ttccaggacc acgctgtatc 240
 tgcaaatgaa cagcctgaga gccgaggaca cggccgtata ttactgtgcg aaagatcttg 300
 gctgggtccga ctcttactac tactactacg gtatggagct ctggggccaa gggaccacgg 360

tcaccgtctc ctcag

375

```

<400> 8
Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser
  1          5          10          15
Leu Arg Leu Ser Cys Thr Ala Ser Gly Phe Thr Phe Ser Ser Tyr Ala
 20          25          30
Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ser
 35          40          45
Ala Ile Ser Gly Ser Gly Gly Thr Thr Phe Tyr Ala Asp Ser Val Lys
 50          55          60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Arg Thr Thr Leu Tyr Leu
 65          70          75          80
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala
 85          90          95
Lys Asp Leu Gly Trp Ser Asp Ser Tyr Tyr Tyr Tyr Tyr Gly Met Asp
100          105          110
Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
115          120

```

<400> 9						
tcctccctgt	ctgcatctgt	aggagacaga	gtcaccttca	cttgccgggc	aagtcaggac	60
attagacgtg	atttaggctg	gtatcagcag	aaaccaggga	aagctcctaa	gcgcctgac	120
tatgctgcat	cccgtttaca	aagtggggtc	ccatcaaggt	tcagcggcag	tggatctggg	180
acagaattca	ctctcacaat	cagcagcctg	cagcctgaag	attttgcaac	ttattactgt	240
ctacagcata	ataattatcc	tcggacgttc	ggccaaggga	ccgaggtgga	aatcatacga	300
ac						302

<400> 10
Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Phe Thr Cys Arg
1 5 10 15

5/31

Ala Ser Gln Asp Ile Arg Arg Asp Leu Gly Trp Tyr Gln Gln Lys Pro
 20 25 30
 Gly Lys Ala Pro Lys Arg Leu Ile Tyr Ala Ala Ser Arg Leu Gln Ser
 35 40 45
 Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr
 50 55 60
 Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys
 65 70 75 80
 Leu Gln His Asn Asn Tyr Pro Arg Thr Phe Gly Gln Gly Thr Glu Val
 85 90 95
 Glu Ile Ile Arg
 100

<210> 11
 <211> 338
 <212> DNA
 <213> Homo sapiens

<400> 11
 gggcccagga ctggtgaagc cttcggagac cctgtccctc acctgcactg tctctgggtg 60
 ctccatcagt aattactact ggagctggat ccggcagccc gccgggaagg gactggagt 120
 gattgggcgt atctatacca gtgggagccc caactacaac ccctccctca agagtcgagt 180
 caccatgtca gtagacacgt ccaagaacca gttctccctg aagctgaact ctgtgaccgc 240
 cgcggacacg gccgtgtatt actgtgcggg aacgattttt ggagtggtta ttatctttga 300
 ctactggggc cagggaaccc tggtcaccgt ctcctcag 338

<210> 12
 <211> 112
 <212> PRT
 <213> Homo sapiens

<400> 12
 Gly Pro Gly Leu Val Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys Thr
 1 5 10 15
 Val Ser Gly Gly Ser Ile Ser Asn Tyr Tyr Trp Ser Trp Ile Arg Gln
 20 25 30
 Pro Ala Gly Lys Gly Leu Glu Trp Ile Gly Arg Ile Tyr Thr Ser Gly
 35 40 45
 Ser Pro Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Met Ser Val
 50 55 60
 Asp Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Asn Ser Val Thr Ala
 65 70 75 80
 Ala Asp Thr Ala Val Tyr Tyr Cys Ala Val Thr Ile Phe Gly Val Val
 85 90 95

6/31

Ile Ile Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 100 105 110

<210> 13
 <211> 322
 <212> DNA
 <213> Homo sapiens

<400> 13
 gacatccaga tgacccagtc tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60
 atcacttgcc gggcaagtca gggcattaga agtgatttag gctggtttca gcagaaacca 120
 gggaaagccc ctaagcgctt gatctatgct gcatccaaat tacaccgtgg ggtcccatca 180
 aggttcagcg gcagtggatc tgggacagaa ttcactctca caatcagccg cctgcagcct 240
 gaagattttg caacttatta ctgtctacag cataatagtt accctctcac ttcggcgga 300
 gggaccaagg tggagatcaa ac 322

<210> 14
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 14
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Ser Asp
 20 25 30
 Leu Gly Trp Phe Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
 35 40 45
 Tyr Ala Ala Ser Lys Leu His Arg Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Arg Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Tyr Pro Leu
 85 90 95
 Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 15
 <211> 376
 <212> DNA
 <213> Homo sapiens

<400> 15
 gaggtgcagc tgttggagtc tgggggaggc ttggtacagc ctgggggggtc cctgagactc 60
 tcctgtgcag cctctggatt caccttttagc agctatgcc a tgagctgggt ccgccaggct 120
 ccagggaagg ggctggagt ggtctcagct attagtggta gtgggtggtat cacatactac 180
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cagcgtgtat 240
 ctgcaaatga acagcctgag agccgaggac acggccgtat attactgtgc gaaagatctg 300
 ggctacgggtg acttttacta ctactactac ggtatggacg tctggggcca agggaccacg 360

7/31

gtcaccgtct cctcag

376

<210> 16
 <211> 125
 <212> PRT
 <213> Homo sapiens

<400> 16
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Ser Gly Ser Gly Gly Ile Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Asp Leu Gly Tyr Gly Asp Phe Tyr Tyr Tyr Tyr Tyr Gly Met
 100 105 110
 Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 115 120 125

<210> 17
 <211> 279
 <212> DNA
 <213> Homo sapiens

<400> 17
 caggagacag agtcaccatc acttgccggg caagtcagag cattagtacc tttttaaatt 60
 ggtatcagca gaaaccaggg aaagccccta aactcctgat ccatgttgca tccagtttac 120
 aagggtggggt cccatcaagg ttcagtggca gtggatctgg gacagatttc actctcacca 180
 tcagcagtct gcaacctgaa gattttgcaa cttactactg tcaacagagt tacaatgccc 240
 cactcacttt cggcggaggg accaaggtgg agatcaaac 279

<210> 18
 <211> 92
 <212> PRT
 <213> Homo sapiens

<400> 18
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Thr
 1 5 10 15
 Phe Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu
 20 25 30

8/31

Ile His Val Ala Ser Ser Leu Gln Gly Gly Val Pro Ser Arg Phe Ser
 35 40 45

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln
 50 55 60

Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Asn Ala Pro
 65 70 75 80

Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 85 90

<210> 19
 <211> 341
 <212> DNA
 <213> Homo sapiens

<400> 19
 cccaggactg gtgaagcctt cggagaccct gtcctcacc tgcactgtct ctggtggctc 60
 catcagtagt tactactgga gttggatccg gcagccccc ggaaggac tggagtggat 120
 tgggtatata tattacagtg ggagcaccaa ctacaacccc tccctcaaga gtcgagtcac 180
 catatcagta gacacgtcca agaaccagtt ctccctgaag ctgagttctg tgaccgctgc 240
 ggacacggcc gtgtattact gtgccaggac gtatagcagt tcgttctact actacggtat 300
 ggacgtctgg ggccaaggga ccacgggtcac cgtctcctca g 341

<210> 20
 <211> 113
 <212> PRT
 <213> Homo sapiens

<400> 20
 Pro Gly Leu Val Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys Thr Val
 1 5 10 15

Ser Gly Gly Ser Ile Ser Ser Tyr Tyr Trp Ser Trp Ile Arg Gln Pro
 20 25 30

Pro Gly Lys Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser
 35 40 45

Thr Asn Tyr Asn Pro Ser Leu Lys Ser Arg Val Thr Ile Ser Val Asp
 50 55 60

Thr Ser Lys Asn Gln Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala
 65 70 75 80

Asp Thr Ala Val Tyr Tyr Cys Ala Arg Thr Tyr Ser Ser Ser Phe Tyr
 85 90 95

Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 100 105 110

Ser

9/31

<210> 21
 <211> 274
 <212> DNA
 <213> Homo sapiens

<220>
 <221> modified_base
 <222> (240)
 <223> a, c, t, g, other or unknown

<400> 21
 agagccaccc tctcctgtag ggccagtcag agtggttcgag gcaggtactt agcctggtag 60
 cagcagaaac ctggccaggc tcccaggctc ctcactatg gtgcatccag cagggccact 120
 ggcattccag acaggttcag tggcagtgag tctgggacag acttcactct caccatcagc 180
 agactggagc ctgaagattt tgcagtgttt tactgtcagc agtatggtag ttcacctcgn 240
 acgttcggcc aagggaacaa ggtggaaatc aaac 274

<210> 22
 <211> 91
 <212> PRT
 <213> Homo sapiens

<400> 22
 Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Arg Gly Arg Tyr
 1 5 10 15
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 20 25 30
 Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser Gly
 35 40 45
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu Pro
 50 55 60
 Glu Asp Phe Ala Val Phe Tyr Cys Gln Gln Tyr Gly Ser Ser Pro Arg
 65 70 75 80
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 85 90

<210> 23
 <211> 367
 <212> DNA
 <213> Homo sapiens

<400> 23
 gaggtgcagc tggttgagtc tgggggaggc ttggtacagc ctgggggggc cctgagactc 60
 tcctgtgcag cctctggatt cacctttagc agctatgcca tgagctgggt ccgccaggct 120
 ccaggaagg ggctggagtg ggtctcaggt attactggga gtggtggtag tacatactac 180
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cagcgtgtat 240
 ctgcaaatga acagcctgag agccgaggac acggccgtat attactgtgc gaaagatcca 300
 gggactacgg tgattatgag ttggttcgac ccctggggcc agggaaccct ggtcaccgtc 360
 tcctcag 367

10/31

<210> 24
 <211> 122
 <212> PRT
 <213> Homo sapiens

<400> 24
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Gly Ile Thr Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Asp Pro Gly Thr Thr Val Ile Met Ser Trp Phe Asp Pro Trp
 100 105 110
 Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 25
 <211> 320
 <212> DNA
 <213> Homo sapiens

<400> 25
 gaactgtggc tgcaccatct gtcttcatct tcccgccatc tgatgagcag ttgaaatctg 60
 gaactgcctc tggtgtgtgc ctgctgaata acttctatcc cagagaggcc aaagtacagt 120
 ggaaggtgga taacgccctc caatcgggta actcccagga gagtgtcaca gagcaggaca 180
 gcaaggacag cacctacagc ctacgcagca ccctgacgct gagcaaagca gactacgaga 240
 aacacaaagt ctacgcctgc gaagtcaccc atcagggcct gagctcgccc gtcacaaaga 300
 gcttcaacag gggagagtgt 320

<210> 26
 <211> 106
 <212> PRT
 <213> Homo sapiens

<400> 26
 Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 1 5 10 15
 Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 20 25 30

11/31

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 35 40 45

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 50 55 60

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 65 70 75 80

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 85 90 95

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 100 105

<210> 27
 <211> 978
 <212> DNA
 <213> Homo sapiens

<400> 27
 gcctccacca agggcccatc ggtcttcccc ctggcgccct gctccaggag cacctccgag 60
 agcacagcgg ccctgggctg cctgggtcaag gactacttcc ccgaaccggg gacgggtgtcg 120
 tggaaactcag gcgctctgac cagcggcggtg cacaccttcc cagctgtcct acagtcctca 180
 ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcaacttcgg caccagacc 240
 tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc 300
 aaatgttggtg tcgagtgcc accgtgcccc gcaccacctg tggcaggacc gtcagtcttc 360
 ctcttcccc caaaacccaa ggacaccctc atgatctccc ggaccctga ggtcacgtgc 420
 gtgggtggtg acgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc 480
 gtggaggtgc ataatgcca gacaaagcca cgggaggagc agttcaacag cacgttccgt 540
 gtggtcagcg tcctcaccgt tgtgcaccag gactggctga acggcaagga gtacaagtgc 600
 aaggtctcca acaaaggcct cccagcccc atcgagaaaa ccatctccaa aaccaaggg 660
 cagccccgag aaccacaggt gtacaccctg ccccatccc gggaggagat gaccaagaac 720
 caggtcagcc tgacctgcct ggtcaaaggc ttctaccca gcgacatcgc cgtggagtgg 780
 gagagcaatg ggcagccgga gaacaactac aagaccacac ctcccatgct ggactccgac 840
 ggctccttct tcctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac 900
 gtcttctcat gtcctgtgat gcatgaggct ctgcacaacc actacacgca gaagagcctc 960
 tccctgtctc cgggtaaa 978

<210> 28
 <211> 326
 <212> PRT
 <213> Homo sapiens

<400> 28
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

12/31

```

Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr
 65                               70                               75                               80

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
                               85                               90                               95

Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro
                               100                              105                              110

Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
                               115                              120                              125

Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 130                              135                              140

Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
 145                              150                              155                              160

Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn
                               165                              170                              175

Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
                               180                              185                              190

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
 195                              200                              205

Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
 210                              215                              220

Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
 225                              230                              235                              240

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
                               245                              250                              255

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
                               260                              265                              270

Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
 275                              280                              285

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
 290                              295                              300

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 305                              310                              315                              320

Ser Leu Ser Pro Gly Lys
                               325

```

<210> 29

<211> 296

<212> DNA

<213> Homo sapiens

13/31

<400> 29
 caggtgcagc tgggtggagtc tggggggaggc ttggtcaagc ctggaggggtc cctgagactc 60
 tcctgtgcag cctctggatt caccttcagt gactactaca tgagctggat ccgccaggct 120
 ccagggaagg ggctggagtg ggtttcatac attagtagta gtggtagtag catatactac 180
 gcagactctg tgaagggccg attcaccatc tccagggaca acgccaagaa ctcactgtat 240
 ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgc gagaga 296

<210> 30
 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 30
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30
 Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Tyr Ile Ser Ser Ser Gly Ser Thr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

<210> 31
 <211> 296
 <212> DNA
 <213> Homo sapiens

<400> 31
 gaggtgcagc tggtggagtc tggggggaggc ttggtacagc ctgggggggtc cctgagactc 60
 tcctgtgcag cctctggatt caccttttagc agctatgccca tgagctgggt ccgccaggct 120
 ccagggaagg ggctggagtg ggtctcagct attagtggta gtggtggtag cacatactac 180
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240
 ctgcaaatga acagcctgag agccgaggac acggccgtat attactgtgc gaaaga 296

<210> 32
 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 32
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

14/31

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys

<210> 33
 <211> 296
 <212> DNA
 <213> Homo sapiens

<400> 33
 caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggggac cctgtccctc 60
 acctgcgctg tctctggtgg ctccatcagc agtagtaact ggtggagttg ggtccgccag 120
 cccccaggga aggggctgga gtggattggg gaaatctatc atagtgggag caccaactac 180
 aaccgcgtccc tcaagagtcg agtcaccata tcagtagaca agtccaagaa ccagttctcc 240
 ctgaagctga gctctgtgac cgccgcggac acggccgtgt attactgtgc gagaga 296

<210> 34
 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 34
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gly
 1 5 10 15

Thr Leu Ser Leu Thr Cys Ala Val Ser Gly Gly Ser Ile Ser Ser Ser
 20 25 30

Asn Trp Trp Ser Trp Val Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp
 35 40 45

Ile Gly Glu Ile Tyr His Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu
 50 55 60

Lys Ser Arg Val Thr Ile Ser Val Asp Lys Ser Lys Asn Gln Phe Ser
 65 70 75 80

Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

15/31

<210> 35
 <211> 293
 <212> DNA
 <213> Homo sapiens

<400> 35
 caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggagac cctgtccctc 60
 acctgcactg tctctggtgg ctccatcagt agttactact ggagctggat ccggcagccc 120
 ccaggaagg gactggagtg gattgggtat atctattaca gtgggagcac caactacaac 180
 ccctccctca agagtcgagt caccatatca gtagacacgt ccaagaacca gttctccctg 240
 aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtgcgag aga 293

<210> 36
 <211> 97
 <212> PRT
 <213> Homo sapiens

<400> 36
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Tyr
 20 25 30
 Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60
 Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80
 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg

<210> 37
 <211> 290
 <212> DNA
 <213> Homo sapiens

<400> 37
 gaaattgtgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagccacc 60
 ctctcctgca gggccagtc gagtgtagc agcagctact tagcctggta ccagcagaaa 120
 cctggccagg ctcccaggct cctcatctat ggtgcatcca gcagggccac tggcatccca 180
 gacaggttca gtggcagtg gtctgggaca gacttcactc tcaccatcag cagactggag 240
 cctgaagatt ttgcagtgt ttactgtcag cagtatggta gctcacctcc 290

<210> 38
 <211> 96
 <212> PRT
 <213> Homo sapiens

16/31

<400> 38

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95

<210> 39

<211> 288

<212> DNA

<213> Homo sapiens

<220>

<221> modified_base

<222> (288)

<223> a, c, t, g, other or unknown

<400> 39

gacatccaga tgaccagtc tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60
 atcacttgcc gggcaagtca gggcattaga aatgathtag gctggatatca gcagaaacca 120
 gggaaagccc ctaagcgct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca 180
 aggttcagcg gcagtggatc tgggacagaa ttcactctca caatcagcag cctgcagcct 240
 gaagattttg caacttatta ctgtctacag cataatagtt accctccn 288

<210> 40

<211> 96

<212> PRT

<213> Homo sapiens

<400> 40

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Asp
 20 25 30
 Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
 35 40 45
 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Tyr Pro Pro
85 90 95

<400>	41						
gacatccaga	tgacccagtc	tccatcctcc	ctgtctgcat	ctgtaggaga	cagagtcacc	60	
atcacttgcc	gggcaagtca	gagcattagc	agctatttaa	attgggtatca	gcagaaacca	120	
gggaaagccc	ctaagctcct	gatctatgct	gcatccagtt	tgcaaagtgg	gggtcccatca	180	
aggttccagt	cagctggatc	tgggacagat	ttcactctca	ccatcagcag	tctgcaacct	240	
gaaqattttt	gcacttacta	ctgtcaacag	agttacagta	cccctcch		288	

<400> 42
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

```
<400> 43
caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggagac cctgtccctc 60
acctgcactg tctctggttg ctccatcagt agttactact ggagctggat ccggcagccc 120
gccgggaagc gactggagtg gattgggcgt atctatacca gtgggagcac caactacaac 180
ccctccctca agatgcagtg caccatgtca gtagacacgt ccaagaacca gttctccctg 240
aagctgagct ctgtgacgc cgcggacacg gccgtgtatt actgtgcgag aga      293
```

8/15/2007, EAST Version: 2.1.0.14

18/31

<212> PRT

<213> Homo sapiens

<400> 44

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Tyr
 20 25 30

Tyr Trp Ser Trp Ile Arg Gln Pro Ala Gly Lys Gly Leu Glu Trp Ile
 35 40 45

Gly Arg Ile Tyr Thr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

Ser Arg Val Thr Met Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg

<210> 45

<211> 470

<212> PRT

<213> Homo sapiens

<400> 45

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Thr Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Ser Tyr Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Ala Ile Ser Gly Ser Gly Gly Thr Thr Phe Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Arg Thr
 85 90 95

Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Lys Asp Leu Gly Trp Ser Asp Ser Tyr Tyr Tyr Tyr
 115 120 125

Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 130 135 140

19/31

Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg	145	150	155	160
Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	165	170	175	
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	180	185	190	
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	195	200	205	
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Asn	Phe	Gly	Thr	Gln	Thr	210	215	220	
Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	225	230	235	240
Thr	Val	Glu	Arg	Lys	Cys	Cys	Val	Glu	Cys	Pro	Pro	Cys	Pro	Ala	Pro	245	250	255	
Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	260	265	270	
Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	275	280	285	
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	290	295	300	
Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	305	310	315	320
Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Val	His	Gln	Asp	Trp	325	330	335	
Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	340	345	350	
Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Gln	Pro	Arg	Glu	355	360	365	
Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	370	375	380	
Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	385	390	395	400
Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	405	410	415	
Thr	Pro	Pro	Met	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	420	425	430	
Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	435	440	445	

20/31

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 450 455 460

Ser Leu Ser Pro Gly Lys
 465 470

<210> 46

<211> 470

<212> PRT

<213> Homo sapiens

<400> 46

Met Glu Phe Gly Leu Ser Trp Leu Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala
 65 70 75 80

Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn
 85 90 95

Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110

Tyr Tyr Cys Ala Lys Gly Tyr Ser Ser Gly Trp Tyr Tyr Tyr Tyr Tyr
 115 120 125

Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 130 135 140

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 145 150 155 160

Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 165 170 175

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 180 185 190

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 195 200 205

Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr
 210 215 220

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 225 230 235 240

21/31

Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro
 245 250 255
 Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
 260 265 270
 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 275 280 285
 Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
 290 295 300
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn
 305 310 315 320
 Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
 325 330 335
 Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
 340 345 350
 Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
 355 360 365
 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
 370 375 380
 Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
 385 390 395 400
 Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
 405 410 415
 Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
 420 425 430
 Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
 435 440 445
 Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 450 455 460
 Ser Leu Ser Pro Gly Lys
 465 470

<210> 47

<211> 236

<212> PRT

<213> Homo sapiens

<400> 47

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1 5 10 15
 Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Phe Pro Ser Ser
 20 25 30

22/31

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45
 Gln Gly Ile Arg Asn Asp Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60
 Ala Pro Lys Arg Leu Ile Tyr Ala Ala Ser Arg Leu His Arg Gly Val
 65 70 75 80
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr
 85 90 95
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln
 100 105 110
 His Asn Ser Tyr Pro Cys Ser Phe Gly Gln Gly Thr Lys Leu Glu Ile
 115 120 125
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
 130 135 140
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
 145 150 155 160
 Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
 165 170 175
 Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
 180 185 190
 Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
 195 200 205
 Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 210 215 220
 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

<210> 48

<211> 236

<212> PRT

<213> Homo sapiens

<400> 48

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Trp
 1 5 10 15
 Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
 20 25 30
 Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
 35 40 45
 Gln Gly Ile Arg Asn Asp Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys
 50 55 60

23/31

Ala Pro Lys Arg Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
 65 70 75 80
 Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr
 85 90 95
 Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln
 100 105 110
 His Asn Ser Tyr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
 115 120 125
 Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
 130 135 140
 Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
 145 150 155 160
 Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
 165 170 175
 Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
 180 185 190
 Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
 195 200 205
 Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
 210 215 220
 Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235

<210> 49

<211> 470

<212> PRT

<213> Homo sapiens

<400> 49

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Ile Lys Gly
 1 5 10 15
 Val Gln Cys Gln Ala Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Asp Tyr Tyr Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ser Tyr Ile Ser Ser Ser Gly Ser Thr Arg Asp Tyr Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95

24/31

Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Val Arg Asp Gly Val Glu Thr Thr Phe Tyr Tyr Tyr Tyr
 115 120 125
 Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 130 135 140
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 145 150 155 160
 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 165 170 175
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 180 185 190
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 195 200 205
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr
 210 215 220
 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 225 230 235 240
 Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro
 245 250 255
 Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
 260 265 270
 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 275 280 285
 Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
 290 295 300
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn
 305 310 315 320
 Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp
 325 330 335
 Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
 340 345 350
 Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu
 355 360 365
 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
 370 375 380
 Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
 385 390 395 400

[illegible]

```
<210> 50
<211> 473
<212> PRT
<213> Homo sapiens
```

<400> 50																
Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Ile	Ile	Lys	Gly	
1				5					10					15		
Val	Gln	Cys	Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	
			20					25					30			
Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	
		35					40					45				
Ser	Asp	Tyr	Tyr	Met	Ser	Trp	Ile	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
	50					55					60					
Glu	Trp	Val	Ser	Tyr	Ile	Ser	Ser	Ser	Gly	Ser	Thr	Ile	Tyr	Tyr	Ala	
65					70					75					80	
Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	
				85					90					95		
Ser	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	
			100					105					110			
Tyr	Tyr	Cys	Ala	Arg	Val	Leu	Arg	Phe	Leu	Glu	Trp	Leu	Leu	Tyr	Tyr	
		115					120					125				
Tyr	Tyr	Tyr	Tyr	Gly	Met	Asp	Val	Trp	Gly	Gln	Gly	Thr	Thr	Val	Thr	
		130				135					140					
Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	
145					150					155					160	
Cys	Ser	Arg	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	
				165					170					175		
Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	
			180					185					190			

26/31

Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly
 195 200 205
 Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Asn Phe Gly
 210 215 220
 Thr Gln Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys
 225 230 235 240
 Val Asp Lys Thr Val Glu Arg Lys Cys Cys Val Glu Cys Pro Pro Cys
 245 250 255
 Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
 260 265 270
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
 275 280 285
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
 290 295 300
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
 305 310 315 320
 Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val His
 325 330 335
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 340 345 350
 Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln
 355 360 365
 Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met
 370 375 380
 Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
 385 390 395 400
 Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
 405 410 415
 Tyr Lys Thr Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu
 420 425 430
 Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
 435 440 445
 Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
 450 455 460
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 465 470

<210> 51
 <211> 236

27/31

<212> PRT

<213> Homo sapiens

<400> 51

```

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1              5              10              15

Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
      20              25              30

Leu Ser Ala Ser Val Gly Asp Arg Val Thr Phe Thr Cys Arg Ala Ser
      35              40              45

Gln Asp Ile Arg Arg Asp Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys
 50              55              60

Ala Pro Lys Arg Leu Ile Tyr Ala Ala Ser Arg Leu Gln Ser Gly Val
 65              70              75              80

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr
      85              90              95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln
      100              105              110

His Asn Asn Tyr Pro Arg Thr Phe Gly Gln Gly Thr Glu Val Glu Ile
      115              120              125

Ile Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
      130              135              140

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
      145              150              155              160
      ^
Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
      165              170              175

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
      180              185              190

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
      195              200              205

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
      210              215              220

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
      225              230              235

```

<210> 52

<211> 236

<212> PRT

<213> Homo sapiens

<400> 52

```

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
 1              5              10              15

```

28/31

```

Phe Pro Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
      20              25              30
Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
      35              40              45
Gln Gly Ile Arg Asn Asp Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys
      50              55              60
Ala Pro Lys Arg Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val
      65              70              75              80
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr
      85              90              95
Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln
      100             105             110
His Asn Ser Tyr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
      115             120             125
Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
      130             135             140
Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
      145             150             155             160
Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
      165             170             175
Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
      180             185             190
Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
      195             200             205
Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
      210             215             220
Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
      225             230             235

```

<210> 53

<211> 326

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Consensus
sequence

<220>

<221> modified_base

<222> (289)

<223> a, c, t, g, other or unknown

29/31

```

<400> 53
gacatccaga tgacccagty tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60
wtcacttgcc gggcaagtca ggrcattaga mrtgatttag gctggwtca gcagaaacca 120
gggaaagcyc ctaagcgctt gatctatgct gcatccmrwt trcammgwgg ggtcccatca 180
aggttcagcg gcagtggatc tgggacagaa ttcactctca caatcagcmg cctgcagcct 240
gaagattttg caacttatta ctgtytacar cataatartr aycckybns ktttygcsrr 300
gggaccrags tggaratcaw acgaac                                     326

```

```

<210> 54
<211> 322
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: Consensus
sequence

```

```

<400> 54
gacatccaga tgacccagtc tccatcctcc ctgtctgcat ctgyaggaga cagagtcacc 60
atcacttgcc gggcaagtca gagcattagy asctwttaa attgggtatca gcagaaacca 120
gggaaagccc ctaarctcct gatcyatgyt gcatccagtt trcaargtgg ggtcccatca 180
aggttcagtg gcagtggatc tgggacagat ttcactctca ccatcagcag tctgcaacct 240
gaagattttg caacttacta ctgtcaacag agttacartr cccayyhc tttcggcgga 300
gggaccaagg tggagatcaa ac                                     322

```

```

<210> 55
<211> 325
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: Consensus
sequence

```

```

<220>
<221> modified_base
<222> (291)
<223> a, c, t, g, other or unknown

```

```

<400> 55
gaaatttgtt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagccacc 60
ctctcctgya gggccagtcg gactgttmgc rgcagstact tagcctggta ccagcagaaa 120
cctggccagg ctcccaggct cctcatctat ggtgcatcca gcagggccac tggcatccca 180
gacaggttca gtggcagtggt gtctgggaca gacttcactc tcaccatcag cagactggag 240
cctgaagatt ttgcagtgtw ttactgtcag cagtatggta gytcacctcs nacgttcggc 300
caagggacca aggtggaaat caaac                                     325

```

```

<210> 56
<211> 376
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: Consensus
sequence

```


30/31

```

<400> 56
caggtgcagc tggaggagtc tgggggagggc ttggtcaagc ctggagggtc cctgagactc 60
tcctgtgcag cctctggatt cacyttcagt gactactaya tgagctggat ccgccaggct 120
ccagggaagg ggctggartg ggtttcatac attagtagta gtggtagtag cakakactac 180
gcagactctg tgaagggccc attcaccatc tccaggagaca acgccaagaa ctcactgtat 240
ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgy gagagatgga 300
gtggaaacta ctttttacta ctactactac ggtatggacg tctggggcca agggaccacg 360
gtcaccgtct cctcag 376

```

```

<210> 57
<211> 358
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: Consensus
sequence

```

```

<220>
<221> modified_base
<222> (337)
<223> a, c, t, g, other or unknown

```

```

<400> 57
caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggagac cctgtccctc 60
acctgcactg tctctgggtg ctccatcagt arttactact ggagctggat ccggcagccc 120
gccgggaagg gactggagtg gattgggcgt atctatacca gtgggagcmc caactacaac 180
ccctccctca agagtgcagt caccatgtca gtagacacgt ccaagaacca gttctccctg 240
aagctgarct ctgtgaccgc cgcggacacg gccgtgtatt actgtgcggt aacgattttt 300
ggagtgggta ttatctttga ctactggggc cagrganccc tggtcaccgt ctcctcag 358

```

```

<210> 58
<211> 418
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: Consensus
sequence

```

```

<400> 58
caggtgcagc tgttggagtc tgggggagggc ttggtacagc ctgggggggtc cctgagactc 60
tcctgtgcag cctctggatt caccttttagc agctatgcc a tgactgggt ccgccaggct 120
ccagggaagg ggctggagtg ggtctcagst attastggka gtggtggtab yacatwctac 180
gcagactccg tgaagggccc gttcaccatc tccagagaca attccargam cacgctgtat 240
ctgcaaatga acagcctgag agccgaggac acggccgtat attactgtgc gaaagatctk 300
ggctrsksyg actyttacta ctactactac ggtatggacg tctggggcca agggacyacg 360
gtgattatga gttggttcga ccctggggc cagggaaccc tggtcaccgt ctcctcag 418

```

```

<210> 59
<211> 364
<212> DNA
<213> Artificial Sequence

```

31/31

<220>

<223> Description of Artificial Sequence: Consensus
sequence

<400> 59

```
caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggagac cctgtccctc 60
acctgcactg tctctggtgg ctccatcagt agttactact ggagytggat ccggcagccc 120
ccagggaagg gactggagtg gattgggtat atctattaca gtgggagcac caactacaac 180
ccctccctca agagtcgact caccatatca gtagacacgt ccaagaacca gttctccctg 240
aagctgagyt ctgtgaccgc tgcggacacg gccgtgtatt actgtgccag gacgtatagc 300
agttcgttct actactacgg tatggacgct tggggccaag ggaccacggt caccgtctcc 360
tcag                                     364
```

<210> 60

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Gly-Ser Linker

<400> 60

```
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
  1               5             10           15
```

Applicants : PFIZER INC., et al.
Filed : 20 December 2001 (20.12.01)
Title : ANTIBODIES TO INSULIN-LIKE GROWTH
FACTOR I RECEPTOR
Agent's Ref. : ABX-PF/2 PCT

Indications Relating to the Expert Solution in
Respect of Deposited Biological Material
Referred to in the Description

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 4.9.2 in the description on

page 75, lines 8-18

The deposit was made in:

American Type Culture Collection (ATCC)
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

on December 12, 2000, under Accession Number PTA 2789.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by

- 2 -

the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

- 3 -

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.

Applicants : PFIZER INC., et al.
Filed : 20 December 2001 (20.12.01)
Title : ANTIBODIES TO INSULIN-LIKE GROWTH
FACTOR I RECEPTOR
Agent's Ref. : ABX-PF/2 PCT

Indications Relating to the Expert Solution in
Respect of Deposited Biological Material
Referred to in the Description

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 4.17.3 in the description on

page 75, lines 8-18

The deposit was made in:

American Type Culture Collection (ATCC)
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

on December 12, 2000, under Accession Number PTA 2793.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by

- 2 -

the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

- 3 -

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.

Applicants : PFIZER INC., et al.
Filed : 20 December 2001 (20.12.01)
Title : ANTIBODIES TO INSULIN-LIKE GROWTH
FACTOR I RECEPTOR
Agent's Ref. : ABX-PF/2 PCT

Indications Relating to the Expert Solution in
Respect of Deposited Biological Material
Referred to in the Description

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 2.12.1 in the description on

page 75, lines 8-18

The deposit was made in:

American Type Culture Collection (ATCC)
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

on December 12, 2000, under Accession Number PTA 2792.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by

- 2 -

the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

- 3 -

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.

Applicants : PFIZER INC., et al.
Filed : 20 December 2001 (20.12.01)
Title : ANTIBODIES TO INSULIN-LIKE GROWTH
FACTOR I RECEPTOR
Agent's Ref. : ABX-PF/2 PCT

Indications Relating to the Expert Solution in
Respect of Deposited Biological Material
Referred to in the Description

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 2.13.2 in the description on

page 75, lines 8-18

The deposit was made in:

American Type Culture Collection (ATCC)
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

on December 12, 2000, under Accession Number PTA 2788.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by

- 2 -

the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

- 3 -

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.

Applicants : PFIZER INC., et al.
Filed : 20 December 2001 (20.12.01)
Title : ANTIBODIES TO INSULIN-LIKE GROWTH
FACTOR I RECEPTOR
Agent's Ref. : ABX-PF/2 PCT

Indications Relating to the Expert Solution in
Respect of Deposited Biological Material
Referred to in the Description

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 2.14.3 in the description on

page 75, lines 8-18

The deposit was made in:

American Type Culture Collection (ATCC)
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

on December 12, 2000, under Accession Number PTA 2790.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by

- 2 -

the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

- 3 -

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.

Applicants : PFIZER INC., et al.
Filed : 20 December 2001 (20.12.01)
Title : ANTIBODIES TO INSULIN-LIKE GROWTH
FACTOR I RECEPTOR
Agent's Ref. : ABX-PF/2 PCT

Indications Relating to the Expert Solution in
Respect of Deposited Biological Material
Referred to in the Description

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 3.1.1 in the description on

page 75, lines 8-18

The deposit was made in:

American Type Culture Collection (ATCC)
10801 University Boulevard
Manassas, Virginia 20110-2209
United States of America

on December 12, 2000, under Accession Number PTA 2791.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by

- 2 -

the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

- 3 -

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.